



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 5/06, A61K 38/55</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 96/40743</b> <b>(43) International Publication Date:</b> 19 December 1996 (19.12.96)
<b>(21) International Application Number:</b> PCT/US96/09285 <b>(22) International Filing Date:</b> 5 June 1996 (05.06.96)  <b>(30) Priority Data:</b> 08/483,470                      7 June 1995 (07.06.95)                      US  <b>(60) Parent Application or Grant</b> (63) Related by Continuation US    08/483,470 (CIP) Filed on    7 June 1995 (07.06.95)  <b>(71) Applicant (for all designated States except US):</b> COR THER- APEUTICS, INC. [US/US]; 256 East Grand Avenue, South San Francisco, CA 94080 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> MARLOWE, Charles, K. [US/US]; 50 Sequoia Court, San Carlos, CA 94070 (US). SCARBOROUGH, Robert, M. [US/US]; 2544 Belmont Canyon Road, Belmont, CA 94002 (US). LAIBELMAN, Alan, M. [US/US]; 427 Waverley, Menlo Park, CA 94025 (US). SINHA, Uma [IN/US]; 808 Junipero Serra Boulevard, San Francisco, CA 94127 (US). ZHU, Bing-Yan [CA/US]; 3325 Adelaide Way, Belmont, CA 94002-1223 (US).	<b>(74) Agents:</b> MENDLEIN, John, D. et al.; Cooley Godward Castro Huddleson & Tatum, Five Palo Alto Square, 3000 El Camino Real, Palo Alto, CA 94306-2155 (US).  <b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished          upon receipt of that report.</i>	
<b>(54) Title:</b> INHIBITORS OF FACTOR Xa  <b>(57) Abstract</b>  Novel compounds, their salts and compositions related thereto having activity against mammalian factor Xa are disclosed. The compounds are useful <i>in vitro</i> or <i>in vivo</i> for preventing or treating coagulation disorders.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

5

## INHIBITORS OF FACTOR Xa

### Field of the Invention

This invention relates to novel arginine and arginine mimetic-containing  
10 compounds which are potent and highly selective inhibitors of isolated factor Xa or  
when assembled in the prothrombinase complex. In another aspect, the present  
invention relates to novel peptide and peptide mimetic analogs, their  
pharmaceutically acceptable salts, and pharmaceutically acceptable compositions  
thereof which are useful as potent and specific inhibitors of blood coagulation in  
15 mammals. In yet another aspect, the invention relates to methods for using these  
inhibitors as therapeutic agents for disease states in mammals characterized by  
coagulation disorders.

### Background of the Invention

20 Hemostasis, the control of bleeding, occurs by surgical means, or by the  
physiological properties of vasoconstriction and coagulation. This invention is  
particularly concerned with blood coagulation and ways in which it assists in  
maintaining the integrity of mammalian circulation after injury, inflammation, disease,  
congenital defect, dysfunction or other disruption. Under normal hemostatic  
25 circumstances, the body maintains an acute balance of clot formation and clot  
removal (fibrinolysis). The blood coagulation cascade involves the conversion of a  
variety of inactive enzymes (zymogens) into active enzymes which ultimately convert  
the soluble plasma protein fibrinogen into an insoluble matrix of highly cross-linked  
fibrin, Davie, E.J. *et al.*, "The Coagulation Cascade: Initiation, Maintenance and  
30 Regulation", *Biochemistry*, 30, 10363-10370 (1991). These plasma glycoprotein  
zymogens include Factor XII, Factor XI, Factor IX, Factor X, Factor VII, and  
prothrombin. Blood coagulation follows either the intrinsic pathway, where all of the  
protein components are present in blood, or the extrinsic pathway, where the cell-  
membrane protein tissue factor plays a critical role. Clot formation occurs when  
35 fibrinogen is cleaved by thrombin to form fibrin. Blood clots are composed of  
activated platelets and fibrin.

Blood platelets which adhere to damaged blood vessels are activated and  
incorporated into the clot and thus play a major role in the initial formation and  
stabilization of hemostatic "plugs". In certain diseases of the cardiovascular system,

5 deviations from normal hemostasis push the balance of clot formation and clot  
dissolution towards life-threatening thrombus formation when thrombi occlude blood  
flow in coronary vessels (myocardial infarctions) or limb and pulmonary veins (venous  
thrombosis). Although platelets and blood coagulation are both involved in thrombus  
formation, certain components of the coagulation cascade are primarily responsible  
10 for the amplification or acceleration of the processes involved in platelet aggregation  
and fibrin deposition.

Thrombin is a key enzyme in the coagulation cascade as well as in  
hemostasis. Thrombin plays a central role in thrombosis through its ability to  
catalyze the conversion of fibrinogen into fibrin and through its potent platelet  
15 activation activity. Under normal circumstances, thrombin can also play an  
anticoagulant role in hemostasis through its ability to convert protein C into  
activated protein C (aPC) in a thrombomodulin-dependent manner. However,  
in atherosclerotic arteries these thrombin activities can initiate the formation  
of a thrombus, which is a major factor in pathogenesis of vasoocclusive  
20 conditions such as myocardial infarction, unstable angina, nonhemorrhagic  
stroke and reocclusion of coronary arteries after angioplasty or thrombolytic  
therapy. Thrombin is also a potent inducer of smooth muscle cell  
proliferation and may therefore be involved in a variety of proliferative  
responses such as restenosis after angioplasty and graft induced  
25 atherosclerosis. In addition, thrombin is chemotactic for leukocytes and may  
therefore play a role in inflammation. (Hoover, R.J., et al. Cell, 14, 423  
(1978); Etingin, O.R., et al., Cell, 61, 657 (1990). These observations  
indicate that inhibition of thrombin formation or inhibition of thrombin itself  
may be effective in preventing or treating thrombosis, limiting restenosis and  
30 controlling inflammation.

Direct or indirect inhibition of thrombin activity has been the focus of a variety  
of recent anticoagulant strategies as reviewed by Claeson, G., "Synthetic Peptides  
and Peptidomimetics as Substrates and Inhibitors of Thrombin and Other Proteases  
in the Blood Coagulation System", Blood Coag. Fibrinol. 5, 411-436 (1994). Several  
35 classes of anticoagulants currently used in the clinic directly or indirectly affect  
thrombin (i.e. heparins, low-molecular weight heparins, heparin-like compounds and  
coumarins).

The formation of thrombin is the result of the proteolytic cleavage of its  
precursor prothrombin at the Arg-Thr linkage at positions 271-272 and the Arg-Ile  
40 linkage at positions 320-321. This activation is catalyzed by the prothrombinase

5 complex, which is assembled on the membrane surfaces of platelets, monocytes,  
and endothelial cells. The complex consists of Factor Xa (a serine protease), Factor  
Va (a cofactor), calcium ions and the acidic phospholipid surface. Factor Xa is the  
activated form of its precursor, Factor X, which is secreted by the liver as a 58 kd  
precursor and is converted to the active form, Factor Xa, in both the extrinsic and  
10 intrinsic blood coagulation pathways. Factor X is a member of the calcium ion  
binding, gamma carboxyglutamyl (Gla)-containing, vitamin K dependent, blood  
coagulation glycoprotein family, which also includes Factors VII and IX, prothrombin,  
protein C and protein S (Furie, B., et al., Cell, 53, 505 (1988)). The activity of Factor  
Xa in effecting the conversion of prothrombin to thrombin is dependent on its  
15 inclusion in the prothrombinase complex.

The prothrombinase complex converts the zymogen prothrombin into the  
active procoagulant thrombin. It is therefore understood that Factor Xa catalyzes the  
next-to-last step in the blood coagulation cascade, namely the formation of the  
serine protease thrombin. In turn, thrombin then acts to cleave soluble fibrinogen in  
20 the plasma to form insoluble fibrin.

The location of the prothrombinase complex at the convergence of the  
intrinsic and extrinsic coagulation pathways, and the resulting significant  
amplification of thrombin generation (several hundred-thousand fold faster in  
effecting the conversion of prothrombin to thrombin than Factor Xa in soluble form)  
25 mediated by the complex at a limited number of targeted catalytic units present at  
vascular lesion sites, suggests that inhibition of thrombin generation is a desirable  
method to block uncontrolled procoagulant activity. It has been suggested that  
compounds which selectively inhibit factor Xa may be useful as *in vitro* diagnostic  
agents, or for therapeutic administration in certain thrombotic disorders, see e.g.,  
30 WO 94/13693. Unlike thrombin, which acts on a variety of protein substrates as well  
as at a specific receptor, factor Xa appears to have a single physiologic substrate,  
namely prothrombin.

Plasma contains an endogenous inhibitor of both the factor VIIa-tissue factor  
(TF) complex and factor Xa called tissue factor pathway inhibitor (TFPI). TFPI is a  
35 Kunitz-type protease inhibitor with three tandem Kunitz domains. TFPI inhibits the  
TF/fVIIa complex in a two-step mechanism which includes the initial interaction of the  
second Kunitz domain of TFPI with the active site of factor Xa, thereby inhibiting the  
proteolytic activity of factor Xa. The second step involves the inhibition of the  
TF/fVIIa complex by formation of a quaternary complex TF/fVIIa/TFPI/fXa as  
40 described by Girard, T.J. et al., "Functional Significance of the Kunitz-type Inhibitory

- 5 Domains of Lipoprotein-associated Coagulation Inhibitor", *Nature*, 338, 518-520 (1989).

Polypeptides derived from hematophagous organisms have been reported which are highly potent and specific inhibitors of factor Xa. United States Patent 4,588,587 describes anticoagulant activity in the saliva of the Mexican leech,  
10 *Haementeria officinalis*. A principal component of this saliva was shown to be the polypeptide factor Xa inhibitor, antistasin (ATS), by Nutt, E. *et al.*, "The Amino Acid Sequence of Antistasin, a Potent Inhibitor of Factor Xa Reveals a Repeated Internal Structure", *J. Biol. Chem.*, 263, 10162-10167 (1988).

Another potent and highly specific inhibitor of Factor Xa, called tick  
15 anticoagulant peptide (TAP), has been isolated from the whole body extract of the soft tick *Omithodoros moubata*, as reported by Waxman, L., *et al.*, "Tick Anticoagulant Peptide (TAP) is a Novel Inhibitor of Blood Coagulation Factor Xa" *Science*, 248, 593-596 (1990).

Other polypeptide type inhibitors of factor Xa have been reported including  
20 the following: Condra, C. *et al.*, "Isolation and Structural Characterization of a Potent Inhibitor of Coagulation Factor Xa from the Leech *Haementeria ghilianii*", *Thromb. Haemost.*, 61, 437-441 (1989); Blankenship, D.T. *et al.*, "Amino Acid Sequence of Ghilanten: Anti-coagulant-antimetastatic Principle of the South American Leech, *Haementeria ghilianii*", *Biochem. Biophys. Res. Commun.* 166, 1384-1389 (1990);  
25 Brankamp, R.G. *et al.*, "Ghilantens: Anticoagulants, Antimetastatic Proteins from the South American Leech *Haementeria ghilianii*", *J. Lab. Clin. Med.*, 115, 89-97 (1990); Jacobs, J.W. *et al.*, "Isolation and Characterization of a Coagulation Factor Xa Inhibitor from Black Fly Salivary Glands", *Thromb. Haemost.*, 64, 235-238 (1990); Rigbi, M. *et al.*, "Bovine Factor Xa Inhibiting Factor and Pharmaceutical  
30 Compositions Containing the Same", European Patent Application, 352,903; Cox, A.C., "Coagulation Factor X Inhibitor From the Hundred-pace Snake *Deinagkistrodon acutus*, venom", *Toxicon*, 31, 1445-1457 (1993); Cappello, M. *et al.*, "Ancylostoma Factor Xa Inhibitor: Partial Purification and its Identification as a Major Hookworm-derived Anticoagulant In Vitro", *J. Infect. Dis.*, 167, 1474-1477 (1993); Seymour, J.L.  
35 *et al.*, "Ecotin is a Potent Anticoagulant and Reversible Tight-binding Inhibitor of Factor Xa", *Biochemistry* 33, 3949-3958 (1994).

Factor Xa inhibitory compounds which are not large polypeptide-type inhibitors have also been reported including: Tidwell, R.R. *et al.*, "Strategies for Anticoagulation With Synthetic Protease Inhibitors. Xa Inhibitors Versus Thrombin

- 5     Inhibitors", *Thromb. Res.*, 19, 339-349 (1980); Turner, A.D. *et al.*, "p-Amidino Esters as Irreversible Inhibitors of Factor IXa and Xa and Thrombin", *Biochemistry*, 25, 4929-4935 (1986); Hitomi, Y. *et al.*, "Inhibitory Effect of New Synthetic Protease Inhibitor (FUT-175) on the Coagulation System", *Haemostasis*, 15, 164-168 (1985); Sturzebecher, J. *et al.*, "Synthetic Inhibitors of Bovine Factor Xa and Thrombin. Comparison of Their Anticoagulant Efficiency", *Thromb. Res.*, 54, 245-252 (1989);
- 10     Kam, C.M. *et al.*, "Mechanism Based Isocoumarin Inhibitors for Trypsin and Blood Coagulation Serine Proteases: New Anticoagulants", *Biochemistry*, 27, 2547-2557 (1988); Hauptmann, J. *et al.*, "Comparison of the Anticoagulant and Antithrombotic Effects of Synthetic Thrombin and Factor Xa Inhibitors", *Thromb. Haemost.*, 63, 220-223 (1990); Miyadera, A. *et al.*, Japanese Patent Application JP 6327488;
- 15     Nagahara, T. *et al.*, "Dibasic (Amidinoaryl)propanoic Acid Derivatives as Novel Blood Coagulation Factor Xa Inhibitors", *J. Med. Chem.*, 37, 1200-1207 (1994); Vlasuk, G.P. *et al.*, "Inhibitors of Thrombosis", WO 93/15756; and Brunck, T.K. *et al.*, "Novel Inhibitors of Factor Xa", WO 94/13693.
- 20     A number of inhibitors of trypsin-like enzymes (such as trypsin, enterokinase, thrombin, kallikrein, plasmin, urokinase, plasminogen activators and the like) have been the subject of disclosures. For example, Austen *et al.*, United States Patent 4,593,018 describes oligopeptide aldehydes which are specific inhibitors of enterokinase; Abe *et al.*, United States Patent 5,153,176 describes tripeptide
- 25     aldehydes which have inhibitory activity against multiple serine proteases such as plasmin, thrombin, trypsin, kallikrein, factor Xa, urokinase, etc.; Brunck *et al.*, European Publication WO 93/14779 describes substituted tripeptide aldehydes that are specific inhibitors of trypsin; United States Patents 4,316889, United States Patent 4,399,065, United States Patent 4,478,745 all disclose arginine aldehyde
- 30     inhibitors of thrombin; Balasubramanian *et al.*, United States Patent 5,380,713 describes di and tripeptide aldehydes which are useful for anti-trypsin and anti-thrombin activity; Webb *et al.*, United States Patent 5,371,072 describes tripeptide alpha-keto-amide derivatives as inhibitors of thrombosis and thrombin; Gesellchen *et al.*, European Patent Publications 0479489A2 and 0643073 A, describe tripeptide
- 35     thrombin inhibitors; Veber *et al.*, European Publication WO 94/25051 describes 4-cyclohexylamine derivatives which selectively inhibit thrombin over other trypsin-like enzymes; Tapparelli *et al.*, *J. Biol. Chem.* 268, 4734-4741 (1993) describe selective peptide boronic acid derivatives as inhibitors of thrombin.

5           Alternatively, agents which inhibit the vitamin K-dependent carboxylase enzyme, such as coumarin, have been used to treat coagulation disorders.

          There exists a need for effective therapeutic agents for the regulation of coagulation disorders, and for the prevention and treatment of thrombus formation and other pathological processes in the vasculature induced by thrombin such as  
10   restenosis and inflammation.


#### Summary of the Invention

          The present invention relates to novel peptide and peptide mimetic analogs, their pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug  
15   derivatives, and pharmaceutically acceptable compositions thereof which have particular biological properties and are useful as potent and specific inhibitors of blood coagulation in mammals. In another aspect, the invention relates to methods of using these inhibitors as diagnostic reagents or as therapeutic agents for disease states in mammals which have coagulation disorders, such as in the treatment or  
20   prevention of any thrombotically mediated acute coronary or cerebrovascular syndrome, any thrombotic syndrome occurring in the venous system, any coagulopathy, and any thrombotic complications associated with extracorporeal circulation or instrumentation, and for the inhibition of coagulation in biological samples.

25           In certain embodiments, this invention relates to novel arginine and arginine mimetic-containing compounds which are potent and highly selective inhibitors of isolated factor Xa when assembled in the prothrombinase complex. These compounds show selectivity for factor Xa versus other proteases of the coagulation cascade (e.g. thrombin, etc.) or the fibrinolytic cascade, and are useful as diagnostic  
30   reagents as well as antithrombotic agents.

          In preferred embodiments, the present invention provides compounds of the formula:





5

wherein:

m = 0,1,2,3,4;

n = 0,1,2,3,4;

p = 0,1,2,3,4;

10

q = 0,1,2,3,4;

Y = CHO, COCF<sub>3</sub>, COCF<sub>2</sub>CF<sub>3</sub>, COCO<sub>2</sub>R<sub>7</sub>, COCONR<sub>8</sub>R<sub>9</sub>, B(OR<sub>10</sub>)<sub>2</sub>;

where: R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub>, R<sub>10</sub> are the same or different and = H, C<sub>1-6</sub>alkyl,  
C<sub>3-8</sub>cycloalkyl, arylC<sub>1-3</sub>alkyl, heteroarylC<sub>1-3</sub>alkyl;

A = piperidinyl, pyrrolidinyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,  
phenyl, C<sub>3-6</sub>heteroaryl, or is absent;

15

R<sub>1</sub> = H or C<sub>1-3</sub>alkyl;J = O or H<sub>2</sub>;R<sub>2</sub> = H or C<sub>1-3</sub>alkyl;D = N, CH, NCH<sub>2</sub>, NCH<sub>2</sub>CH<sub>2</sub>, CHCH<sub>2</sub>;

20

R<sub>3</sub> = H or C<sub>1-3</sub>alkyl;E = O or H<sub>2</sub>;R<sub>4</sub> = H or CH<sub>3</sub>;

Q = piperidinyl, pyrrolidinyl, C<sub>3-8</sub> cycloalkyl, phenyl, substituted phenyl,  
naphthyl, pyridyl, or is absent;

25

G = N, CH, or is H;

R<sub>5</sub> = H or C<sub>1-3</sub> alkyl, or is absent if G is H;R<sub>6</sub> = H or CH<sub>3</sub>

W = H, arylacyl, heteroarylacyl, arylC<sub>1-3</sub>alkylsulfonyl, arylsulfonyl,  
substituted arylsulfonyl, arylC<sub>1-4</sub>alkenylsulfonyl, C<sub>1-8</sub> alkylsulfonyl,

5 heteroarylC<sub>1-3</sub>alkylsulfonyl, heteroarylsulfonyl, aryloxycarbonyl, C<sub>1-6</sub> alkyloxycarbonyl, arylC<sub>1-3</sub>alkyloxycarbonyl, arylaminocarbonyl, C<sub>1-6</sub>alkylaminocarbonyl, arylC<sub>1-3</sub>alkylaminocarbonyl, HOOC-C<sub>0-3</sub>alkylcarbonyl, or is absent if G is H;

10 X = H, C<sub>1-3</sub>alkyl, NR'R'', NH-C(NR'R'')=NH, NH-C(NHR')=NR'', S-C(NR'R'')=NH, S-C(NHR')=NR'', C(NR'R'')=NH, C(NHR')=NR'', or CR'=NR''; where: R', R'' are the same or different and = H, C<sub>1-6</sub>alkyl, C<sub>1-3</sub>arylalkyl, aryl or where R'R'' forms a cyclic ring containing (CH<sub>2</sub>)<sub>p</sub> where p=2-5, with the proviso that when X is H or C<sub>1-3</sub>alkyl, then A must contain at least one N atom;

15 Z = H, C<sub>1-3</sub>alkyl, NR'R'', NH-C(NR'R'')=NH, NH-C(NHR')=NR'', S-C(NR'R'')=NH, S-C(NHR')=NR'', C(NR'R'')=NH, C(NHR')=NR'', or CR'=NR''; where: R', R'' are the same or different and = H, C<sub>1-6</sub>alkyl, C<sub>1-3</sub>arylalkyl, aryl or where R'R'' forms a cyclic ring containing (CH<sub>2</sub>)<sub>p</sub> where p=2-5, with the proviso that when Z is H or C<sub>1-3</sub>alkyl, then Q must contain at least one N atom;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

In certain aspects of this invention, compounds are provided which are useful as diagnostic reagents. In another aspect, the present invention includes  
 25 pharmaceutical compositions comprising a pharmaceutically effective amount of the compounds of this invention and a pharmaceutically acceptable carrier. In yet another aspect, the present invention includes methods comprising using the above compounds and pharmaceutical compositions for preventing or treating disease states characterized by disorders of the blood coagulation process in mammals, or  
 30 for preventing coagulation in stored blood products and samples. Optionally, the methods of this invention comprise administering the pharmaceutical composition in combination with an additional therapeutic agent such as an antithrombotic and/or a thrombolytic agent and/or an anticoagulant.

The preferred compounds also include their pharmaceutically acceptable  
 35 isomers, hydrates, solvates, salts and prodrug derivatives.

#### Detailed Description of the Invention

## 5 Definitions

In accordance with the present invention and as used herein, the following terms are defined with the following meanings, unless explicitly stated otherwise.

The term "alkenyl" refers to a trivalent straight chain or branched chain unsaturated aliphatic radical.

10 The term "alkyl" refers to saturated aliphatic groups including straight-chain, branched-chain and cyclic groups having the number of carbon atoms specified, or if no number is specified, having up to 12 carbon atoms. The term "cycloalkyl" as used herein refers to a mono-, bi-, or tricyclic aliphatic ring having 3 to 14 carbon atoms and preferably 3 to 7 carbon atoms.

15 The term "aryl" refers to an unsubstituted or substituted aromatic ring, substituted with one, two or three substituents selected from loweralkoxy, loweralkyl, loweralkylamino, hydroxy, halogen, cyano, hydroxyl, mercapto, nitro, thioalkoxy, carboxaldehyde, carboxyl, carboalkoxy and carboxamide, including but not limited to carbocyclic aryl, heterocyclic aryl, and biaryl groups and the like, all of which may be  
20 optionally substituted. Preferred aryl groups include phenyl, halophenyl, loweralkylphenyl, naphthyl, biphenyl, phenanthrenyl, naphthacenyl, and aromatic heterocyclics. The term "heteroaryl" as used herein refers to any aryl group, containing from one to four heteroatoms, selected from the group consisting of nitrogen, oxygen and sulfur.

25 The term "arylalkyl" refers to one, two, or three aryl groups having the number of carbon atoms designated, appended to an alkyl group having the number of carbon atoms designated. Suitable arylalkyl groups include, but are not limited to, benzyl, picolyl, naphthylmethyl, phenethyl, benzyhydryl, trityl, and the like, all of which may be optionally substituted.

30 The terms "halo" or "halogen" as used herein refer to Cl, Br, F or I substituents.

The term "methylene" refers to  $-CH_2-$ .

The term "pharmaceutically acceptable salts" includes salts of compounds derived from the combination of a compound and an organic or inorganic acid.

35 These compounds are useful in both free base and salt form. In practice, the use of the salt form amounts to use of the base form; both acid and base addition salts are within the scope of the present invention.

"Pharmaceutically acceptable acid addition salt" refers to those salts which retain the biological effectiveness and properties of the free bases and which are not  
40 biologically or otherwise undesirable, formed with inorganic acids such as

5 hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

10 "Pharmaceutically acceptable base addition salts" include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Particularly preferred are the ammonium, potassium, sodium, calcium and magnesium salts. Salts derived from pharmaceutically acceptable organic nontoxic bases include salts  
15 of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-diethylaminoethanol, trimethamine, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine,  
20 ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins and the like. Particularly preferred organic nontoxic bases are isopropylamine, diethylamine, ethanolamine, trimethamine, dicyclohexylamine, choline, and caffeine.

"Biological property" for the purposes herein means an *in vivo* effector or  
25 antigenic function or activity that is directly or indirectly performed by a compound of this invention. Effector functions include receptor or ligand binding, any enzyme activity or enzyme modulatory activity, any carrier binding activity, any hormonal activity, any activity in promoting or inhibiting adhesion of cells to an extracellular matrix or cell surface molecules, or any structural role. Antigenic functions include  
30 possession of an epitope or antigenic site that is capable of reacting with antibodies raised against it.

The nomenclature used to describe the peptide compounds of the invention follows the conventional practice where the N-terminal amino group is assumed to be to the left and the carboxy group to the right of each amino acid residue in the  
35 peptide. In the formulas representing selected specific embodiments of the present invention, the amino- and carboxy-terminal groups, although often not specifically shown, will be understood to be in the form they would assume at physiological pH values, unless otherwise specified. Thus, the N-terminal  $H^+$  and C-terminal  $O^-$  at physiological pH are understood to be present though not necessarily specified and  
40 shown, either in specific examples or in generic formulas. Free functional groups on

- 5 the side chains of the amino acid residues can also be modified by amidation, acylation or other substitution, which can, for example, change the solubility of the compounds without affecting their activity.

In the peptides described herein, each gene-encoded residue, where appropriate, is represented by a single letter designation, corresponding to the trivial  
10 name of the amino acid, in accordance with the following conventional list:

	One-Letter	Three-letter	
	<u>Amino Acid</u>	<u>Symbol</u>	<u>Symbol</u>
	Alanine	A	Ala
	Arginine	R	Arg
15	Asparagine	N	Asn
	Aspartic acid	D	Asp
	Cysteine	C	Cys
	Glutamine	Q	Gln
	Glutamic acid	E	Glu
20	Glycine	G	Gly
	Histidine	H	His
	Isoleucine	I	Ile
	Leucine	L	Leu
	Lysine K	Lys	
25	Methionine	M	Met
	Phenylalanine	F	Phe
	Proline P	Pro	
	Serine S	Ser	
	Threonine	T	Thr
30	Tryptophan	W	Trp
	Tyrosine	Y	Tyr
	Valine V	Val	

In addition, the following abbreviations are used in this application:

- "Ala" refers to L-Alanine.
- 35 "D-Ala" refers to D-Alanine.
- " $\beta$ -Ala" refers to 3-aminopropanoic acid.
- "Arg" refers to L-Arginine.
- "D-Arg" refers to D-Arginine.
- "Aib" refers to alpha-aminoisobutyric acid.

40

- 5 "Bn" refers to benzyl.  
"t-Boc" refers to t-butoxycarbonyl.  
"BOP" refers to benzotriazol-1-yloxy-tris-(dimethylamino)phosphonium  
hexafluorophosphate.  
"Brine" means an aqueous saturated solution of sodium chloride.
- 10 "Cbz" refers to benzyloxycarbonyl.  
"CDI" refers to carbonyldiimidazole.  
"DCC" refers to dicyclohexylcarbodiimide.  
"DCM" refers to dichloromethane.  
"DCU" refers to dicyclohexylurea.
- 15 "DIEA" refers to diisopropylethylamine.  
"DMF" refers to N,N-dimethylformamide.  
"EtOAc" refers to ethyl acetate.  
"Fm" refers to 9-fluorenylmethyl.  
"Gly" refers to glycine.
- 20 "HOSu" refers to N-hydroxysuccinimide.  
"HATU" refers to O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl-  
uronium  
hexafluorophosphate.  
"HBTU" refers to O-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyl-uronium  
hexafluorophosphate.
- 25 "HOBt" refers to N-hydroxybenzotriazole.  
"IPA" refers to isopropanol.  
"D-Lys" refers to D-Lysine.  
"MeOH" refers to methanol.
- 30 "NaOAc" refers to sodium acetate.  
"NMM" refers to 4-methylmorpholine.  
"2-NaphthoxyAc" refers to 2-Naphthoxyacetyl.  
"Ph" refers to phenyl.  
"D-Pro" refers to D-proline.
- 35 "Pro" refers to L-proline.  
"TEA" refers to triethylamine.  
"TFA" refers to trifluoroacetic acid.

5 "THF" refers to tetrahydrofuran.

"TsOH" refers to p-toluenesulfonic acid.

The amino acids not encoded genetically are abbreviated as described above or have the meanings commonly accepted in the field.

10 In the compounds of this invention, carbon atoms bonded to four non-identical substituents are asymmetric. Accordingly, the compounds may exist as diastereoisomers, enantiomers or mixtures thereof. The syntheses described herein may employ racemates, enantiomers or diastereomers as starting materials or intermediates. Diastereomeric products resulting from such syntheses may be separated by chromatographic or by crystallization methods, or by other methods  
15 known in the art. Likewise, enantiomeric product mixtures may be separated using the same techniques or by other methods known in the art. Each of the asymmetric carbon atoms, when present in the compounds of this invention, may be in one of two configurations (R or S) and both are within the scope of the present invention. In certain specified preferred embodiments of the compounds shown in the present  
20 application, the L-form of any amino acid residue having an optical isomer is intended unless the D-form is expressly indicated. In the processes described above, the final products may, in some cases, contain a small amount of the products having D or L-form residues, however these products do not affect their therapeutic or diagnostic application.

25 The compounds of the invention are peptides or compounds which contain amino acid subunits which are partially defined in terms of amino acid residues of designated classes. Amino acid residues can be generally grouped into four major subclasses as follows:

30 Acidic: The residue has a negative charge due to loss of H ion at physiological pH and the residue is attracted by aqueous solution.

Basic: The residue has a positive charge due to association with H ion at physiological pH and the residue is attracted by aqueous solution.

35 Neutral/nonpolar: The residues are not charged at physiological pH and the residue is repelled by aqueous solution so as to seek the inner positions in the conformation of a peptide in which it is contained when the peptide is in aqueous medium. These residues are also designated "hydrophobic" herein.

40 Neutral/polar: The residues are not charged at physiological pH, but the residue is attracted by aqueous solution so as to seek the outer positions in the conformation of a peptide in which it is contained when the peptide is in aqueous medium.

5           It is understood, of course, that in a statistical collection of individual residue molecules some molecules will be charged, and some not, and there will be an attraction for or repulsion from an aqueous medium to a greater or lesser extent. To fit the definition of "charged," a significant percentage (at least approximately 25%) of the individual molecules are charged at physiological pH. The degree of attraction  
10 or repulsion required for classification as polar or nonpolar is arbitrary and, therefore, amino acids specifically contemplated by the invention have been classified as one or the other. Most amino acids not specifically named can be classified on the basis of known behavior.

Amino acid residues can be further subclassified as cyclic or noncyclic, and  
15 aromatic or nonaromatic, self-explanatory classifications with respect to the side chain substituent groups of the residues, and as small or large. The residue is considered small if it contains a total of 4 carbon atoms or less, inclusive of the carboxyl carbon. Small residues are, of course, always nonaromatic for naturally occurring protein amino acids.

20           For the naturally occurring protein amino acids, subclassification according to the foregoing scheme is as follows.

Acidic: Aspartic acid and Glutamic acid;

Basic/noncyclic: Arginine, Lysine;

Basic/cyclic: Histidine;

25 Neutral/small: Glycine, Serine, Cysteine, Alanine;

Neutral/polar/large/nonaromatic: Threonine, Asparagine, Glutamine;

Neutral/polar/large/aromatic: Tyrosine;

Neutral/nonpolar/large/nonaromatic: Valine, Isoleucine, Leucine, Methionine;

Neutral/nonpolar/large/aromatic: Phenylalanine, and Tryptophan.

30           The gene-encoded secondary amino acid proline, although technically within the group neutral/nonpolar/large/ cyclic and nonaromatic, is a special case due to its known effects on the secondary conformation of peptide chains, and is not, therefore, included in this defined group.

Certain commonly encountered amino acids, which are not encoded by the  
35 genetic code, include, for example, beta-alanine (b-Ala), or other omega-amino acids, such as 2,3-diamino propionic (2,3-Dap), 2,4-diaminobutyric (2,4-Dab), 4-amino butyric (g-Abu) and so forth, alpha-aminoisobutyric acid (Aib), sarcosine (Sar), ornithine (Om), citrulline (Cit), homoarginine (Har), homolysine (homoLys), n-butylamidinoglycine (Bag), 4-guanidinophenylalanine (4-Gpa), 3-  
40 guanidinophenylalanine (3-Gpa), 4-amidinophenylalanine (4-Apa), 3-amidinophenylalanine (3-Apa), 4-aminocyclohexylglycine (4-Acg), 4-



- 5    aminophenylalanine (4-NH<sub>2</sub>-Phe), 3-aminophenylalanine (3-NH<sub>2</sub>-Phe), 3-piperidinyllalanine (3-PIA), 4-piperidinyllalanine (4-PIA), 3-guanidinopiperidinyllalanine (3-GPIA), 4-guanidinopiperidinyllalanine (4-GPIA), 3-piperidinylglycine (3-PIG), 4-piperidinylglycine (4-PIG), 3-guanidinopiperidinylglycine (3-GPIG), and 4-guanidinopiperidinylglycine (4-GPIG). These also fall conveniently into particular  
10    categories.

Based on the above definitions:

Sar, b-Ala, g-Abu, and Aib are neutral/small;

- Orn, Har, homoLys, Bag, 2,3-Dap, 2,4-Dab, 4-Gpa, 3-Gpa, 4-Apa, 3-Apa, 4-Acg, 4-NH<sub>2</sub>-Phe, 3-NH<sub>2</sub>-Phe, 3-PIA, 4-PIA, 3-GPIA, 4-GPIA, 3-PIG, 4-PIG, 3-GPIG  
15    and 4-GPIG are basic;

Cit, is neutral/polar/ large/nonaromatic; and

The various omega-amino acids are classified according to size as neutral/nonpolar/small (β-Ala, i.e., 3-aminopropionic, 4-aminobutyric) or large (all others).


- 20    Amino acid substitutions for those indicated in the structure/formula provided can be included in peptide compounds within the scope of the invention and can be classified within this general scheme according to their structure.

- In all of the peptides of the invention, one or more amide linkages (-CO-NH-) may optionally be replaced with another linkage which is an isostere such as  
25    -CH<sub>2</sub>NH-, -CH<sub>2</sub>S-, -CH<sub>2</sub>CH<sub>2</sub>-, -CH=CH- (cis and trans), -COCH<sub>2</sub>-, -CH(OH)CH<sub>2</sub>- and -CH<sub>2</sub>SO-. This replacement can be made by methods known in the art. The following references describe preparation of peptide analogs which include these alternative-linking moieties: Spatola, A.F., Vega Data (March 1983), Vol. 1, Issue 3, "Peptide Backbone Modifications" (general review); Spatola, A.F., in "Chemistry and  
30    Biochemistry of Amino Acids, Peptides and Proteins," B. Weinstein, eds., Marcel Dekker, New York, p. 267 (1983) (general review); Morley, J.S., Trends Pharm Sci (1980) pp. 463-468 (general review); Hudson, D., et al., Int J Pept Prot Res (1979) 14:177-185 (-CH<sub>2</sub>NH-, -CH<sub>2</sub>CH<sub>2</sub>-); Spatola, A.F., et al., Life Sci (1986) 38:1243-1249 (-CH<sub>2</sub>-S); Hann, M.M., J Chem Soc Perkin Trans I (1982) 307-314 (-CH=CH-,  
35    cis and trans); Almquist, R.G., et al., J Med Chem (1980) 23:1392-1398 (-COCH<sub>2</sub>-); Jennings-White, C., et al., Tetrahedron Lett (1982) 23:2533 (-COCH<sub>2</sub>-); Szelke, M., et al., European Application EP 45665; CA:97:39405 (1982) (-CH(OH)CH<sub>2</sub>-); Holladay, M.W., et al., Tetrahedron Lett (1983) 24:4401-4404 (-C(OH)CH<sub>2</sub>-); and Hruby, V.J., Life Sci (1982) 31:189-199 (-CH<sub>2</sub>-S-).

5

Preferred Embodiments

In preferred embodiments, the present invention provides compounds of the formula:



10 wherein:

m = 0,1,2,3,4;

n = 0,1,2,3,4;

p = 0,1,2,3,4;

q = 0,1,2,3,4;

15 Y = CHO, COCF<sub>3</sub>, COCF<sub>2</sub>CF<sub>3</sub>, COCO<sub>2</sub>R<sub>7</sub>, COCONR<sub>8</sub>R<sub>9</sub>, B(OR<sub>10</sub>)<sub>2</sub>;

where: R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub>, R<sub>10</sub> are the same or different and = H, C<sub>1-6</sub>alkyl, C<sub>3-8</sub>cycloalkyl, arylC<sub>1-3</sub>alkyl, heteroarylC<sub>1-3</sub>alkyl;

A = piperdiny, pyrrolidiny, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, C<sub>3-6</sub>heteroaryl, or is absent;

20 R<sub>1</sub> = H or C<sub>1-3</sub>alkyl;

J = O or H<sub>2</sub>;

R<sub>2</sub> = H or C<sub>1-3</sub>alkyl;

D = N, CH, NCH<sub>2</sub>, NCH<sub>2</sub>CH<sub>2</sub>, CHCH<sub>2</sub>;

R<sub>3</sub> = H or C<sub>1-3</sub>alkyl;

25 E = O or H<sub>2</sub>;

R<sub>4</sub> = H or CH<sub>3</sub>;

Q = piperdiny, pyrrolidiny, C<sub>3-8</sub> cycloalkyl, phenyl, substituted phenyl, naphthyl, pyridyl, or is absent;


G = N, CH, or H;

30 R<sub>5</sub> = H or C<sub>1-3</sub> alkyl, or is absent if G is H;

- 5             $R_6 =$  H or  $CH_3$ ;
- $W =$  H, arylacetyl, heteroarylacetyl, arylC<sub>1-3</sub>alkylsulfonyl, arylsulfonyl, substituted arylsulfonyl, arylC<sub>1-4</sub>alkenylsulfonyl, C<sub>1-8</sub> alkylsulfonyl, heteroarylC<sub>1-3</sub>alkylsulfonyl, heteroarylsulfonyl, aryloxycarbonyl, C<sub>1-6</sub> alkyloxycarbonyl, arylC<sub>1-3</sub>alkyloxycarbonyl, arylaminocarbonyl, C<sub>1-6</sub> alkylaminocarbonyl, arylC<sub>1-3</sub>alkylaminocarbonyl, HOOC-C<sub>0-3</sub>alkylcarbonyl, or is absent if G is H;
- 10            $X =$  H, C<sub>1-3</sub>alkyl, NR'R'', NH-C(NR'R'')=NH, NH-C(NHR')=NR'', S-C(NR'R'')=NH, S-C(NHR')=NR'', C(NR'R'')=NH, C(NHR')=NR'', or CR'=NR''; where: R', R'' are the same or different and = H, C<sub>1-6</sub>alkyl, C<sub>1-3</sub>arylalkyl, aryl or where R'R'' forms a cyclic ring containing (CH<sub>2</sub>)<sub>p</sub> where p=2-5, with the proviso that when X is H or C<sub>1-3</sub>alkyl, then A must contain at least one N atom;
- 15            $Z =$  H, C<sub>1-3</sub>alkyl, NR'R'', NH-C(NR'R'')=NH, NH-C(NHR')=NR'', S-C(NR'R'')=NH, S-C(NHR')=NR'', C(NR'R'')=NH, C(NHR')=NR'', or CR'=NR''; where: R', R'' are the same or different and = H, C<sub>1-6</sub>alkyl, C<sub>1-3</sub>arylalkyl, aryl or where R'R'' forms a cyclic ring containing (CH<sub>2</sub>)<sub>p</sub> where p=2-5, with the proviso that when Z is H or C<sub>1-3</sub>alkyl, then Q must contain at least one N atom;
- 20           and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

25           derivatives thereof.

             More preferably, compounds of the present invention include those of formula:



5

wherein:

m = 0,1,2,3,4;

n = 0,1,2,3,4;

p = 0,1,2,3,4;

10

q = 0,1,2,3,4;

Y = CHO, COCF<sub>3</sub>, COCF<sub>2</sub>CF<sub>3</sub>, COCO<sub>2</sub>R<sub>7</sub>, COCONR<sub>8</sub>R<sub>9</sub>; where:R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub> are the same or different and = H, C<sub>1-6</sub>alkyl, C<sub>3-8</sub>cycloalkyl, arylC<sub>1-3</sub>alkyl, heteroarylC<sub>1-3</sub>alkyl;

15

A = piperdiny, pyrrolidiny, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, C<sub>3-6</sub>heteroaryl or is absent;R<sub>2</sub> = H or C<sub>1-3</sub>alkyl;Q = piperdiny, pyrrolidiny, C<sub>3-8</sub> cycloalkyl, phenyl, substituted phenyl, naphthyl, pyridyl, or is absent;

20

W = H, arylacyl, heteroarylacyl, arylC<sub>1-3</sub>alkylsulfonyl, arylsulfonyl, substituted arylsulfonyl, arylC<sub>1-4</sub>alkenylsulfonyl, C<sub>1-8</sub> alkylsulfonyl, heteroarylC<sub>1-3</sub>alkylsulfonyl, heteroaryl sulfonyl, aryloxycarbonyl, C<sub>1-6</sub> alkyloxycarbonyl, arylC<sub>1-3</sub>alkyloxycarbonyl, arylaminocarbonyl, C<sub>1-6</sub> alkylaminocarbonyl, arylC<sub>1-3</sub>alkylaminocarbonyl, or HOOC-C<sub>0-3</sub>alkylcarbonyl;

25

X = H, C<sub>1-3</sub>alkyl, NR'R'', NH-C(NR'R'')=NH, NH-C(NHR')=NR'', S-C(NR'R'')=NH, S-C(NHR')=NR'', C(NR'R'')=NH, C(NHR')=NR'', or CR'=NR''; where: R', R'' are the same or different and = H, C<sub>1-6</sub>alkyl, C<sub>1-3</sub>arylalkyl, aryl or where R'R'' forms a cyclic ring containing (CH<sub>2</sub>)<sub>p</sub> where p=2-5, with the proviso that when X is H or C<sub>1-3</sub>alkyl, then A must contain at least one N atom;

30

- 5           Z = H, C<sub>1-3</sub>alkyl, NR'R", NH-C(NR'R")=NH, NH-C(NHR')=NR", S-C(NR'R")=NH, S-C(NHR')=NR", C(NR'R")=NH, C(NHR')=NR", or CR'=NR"; where: R',R" are the same or different and = H, C<sub>1-6</sub>alkyl, C<sub>1-3</sub>arylalkyl, aryl or where R'R" forms a cyclic ring containing (CH<sub>2</sub>)<sub>p</sub> where p=2-5, with the proviso that when Z is H or C<sub>1-3</sub>alkyl, then Q
- 10           must contain at least one N atom;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

- A preferred substituent Y is CHO, COCO<sub>2</sub>R<sub>7</sub>, COCONR<sub>8</sub>R<sub>9</sub>, where: R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub>, are the same or different and = H, C<sub>1-6</sub>alkyl, C<sub>1-8</sub>cycloalkyl, arylC<sub>1-3</sub>alkyl,
- 15   heteroarylC<sub>1-3</sub>alkyl.

A preferred substituent A is piperdiny, pyrrolidiny, cyclopentyl, cyclohexyl, phenyl, C<sub>3-6</sub>heteroaryl, or is absent.

A preferred substituent D is N, CH, NCH<sub>2</sub>.

- A preferred substituent Q is piperdiny, pyrrolidiny, C<sub>3-8</sub> cycloalkyl, phenyl,
- 20   substituted phenyl.

A preferred substituent W is arylC<sub>1-3</sub>alkylsulfonyl, arylsulfonyl, substituted arylsulfonyl, arylC<sub>1-4</sub>alkenylsulfonyl, C<sub>1-8</sub> alkylsulfonyl, heteroarylC<sub>1-3</sub>alkylsulfonyl, heteroarylsulfonyl, C<sub>1-6</sub> alkyloxycarbonyl, arylC<sub>1-3</sub>alkyloxycarbonyl.

- A preferred substituent X is NR'R", NH-C(NR'R")=NH, NH-C(NHR')=NR", S-C(NR'R")=NH, S-C(NHR')=NR", C(NR'R")=NH, C(NHR')=NR"; where: R',R" are the same or different and = H, C<sub>1-6</sub>alkyl.
- 25

A preferred substituent Z is NH-C(NR'R")=NH, NH-C(NHR')=NR", S-C(NR'R")=NH, S-C(NHR')=NR", C(NR'R")=NH, C(NHR')=NR"; where: R',R" are the same or different and = H, C<sub>1-6</sub>alkyl.

- 30           Preferred compounds as a whole may be selected from any combination of the formulas presented in this specification with one or more of the preferred groupings of substituents at a particular location.

- 5 Some preferred embodiments of the invention are shown in the following Table 1.

TABLE 1

	STRUCTURE	Inhibitory Activity (IC <sub>50</sub> ) mM		
		Factor Xa	Prothrombinase	Thrombin
10	H-D-Arg-Gly-Arg-H	0.064	0.83	41
	Boc-D-Arg-Gly-Arg-H	0.050	4.0	>100
	HOCCO-D-Arg-Gly-Arg-H	0.056	0.63	95
	HOOCCH <sub>2</sub> CO-D-Arg-Gly-Arg-H	0.098	0.50	>100
	HOOC(CH <sub>2</sub> ) <sub>2</sub> CO-D-Arg-Gly-Arg-H	0.224	2.0	>100
15	PhCH <sub>2</sub> CH <sub>2</sub> CO-D-Arg-Gly-Arg-H	0.200	2.0	75
	PhCH <sub>2</sub> SO <sub>2</sub> -D-Arg-Gly-Arg-H	0.015	0.014	33
	EtOCO-D-Arg-Gly-Arg-H	0.049	0.135	>100
	2-NaphthoxyAc-D-Arg-Gly-Arg-H	0.550	0.687	>100
	Boc-D-Cit-Gly-Arg-H	5.0	44.0	>100
20	Boc-D-Lys-Gly-Arg-H	10.0	68.0	>100
	Boc-D-Har-Gly-Arg-H	0.100	0.173	405
	Boc-D-Har((CH <sub>3</sub> ) <sub>4</sub> )-Gly-Arg-H	3.0	12.0	>100
	Boc-D-Arg-Ala-Arg-H	0.167	0.081	5
	Boc-D-Arg-D-Ala-Arg-H	0.230	0.800	>10
25	Boc-D-Arg-β-Ala-Arg-H	11.0	14.0	>500
	Boc-D-Arg-Aib-Arg-H	0.552	2.0	>500
	Boc-D-Arg-Gly-Arg-CONHEtPh	0.016	0.023	4

- Other preferred embodiments of the invention are shown but are not limited to the following list of compounds, which have the general structure :

W - (Basic amino acid) - (Neutral/small amino acid) - (Arg or Basic amino acid) - Y

Boc-D-(2,3-Dap)-Gly-Arg-H

Boc-D-(2,4-Dab)-Gly-Arg-H

γ-Abu-Gly-Arg-H

- 35 Boc-D-Om-Gly-Arg-H

Boc-D-homoLys-Gly-Arg-H

Boc-Bag-Gly-Arg-H

Boc-D-4-Gpa-Gly-Arg-H

Boc-D-3-Gpa-Gly-Arg-H

5	Boc-D-4-Apa-Gly-Arg-H Boc-D-3-Apa-Gly-Arg-H Boc-D-4-Acg-Gly-Arg-H Boc-D-(4-NH <sub>2</sub> Phe)-Gly-Arg-H Boc-D-(3-NH <sub>2</sub> Phe)-Gly-Arg-H
10	BnSO <sub>2</sub> -D-(2,3-Dap)-Gly-Arg-H BnSO <sub>2</sub> -D-(2,4-Dab)-Gly-Arg-H BnSO <sub>2</sub> -D-Om-Gly-Arg-H BnSO <sub>2</sub> -D-homoLys-Gly-Arg-H BnSO <sub>2</sub> -Bag-Gly-Arg-H
15	BnSO <sub>2</sub> -D-4-Gpa-Gly-Arg-H BnSO <sub>2</sub> -D-3-Gpa-Gly-Arg-H BnSO <sub>2</sub> -D-4-Apa-Gly-Arg-H BnSO <sub>2</sub> -D-3-Apa-Gly-Arg-H BnSO <sub>2</sub> -D-4-Acg-Gly-Arg-H
20	BnSO <sub>2</sub> -D-(4-NH <sub>2</sub> Phe)-Gly-Arg-H BnSO <sub>2</sub> -D-(3-NH <sub>2</sub> Phe)-Gly-Arg-H BnSO <sub>2</sub> -D-(2,3-Dap)-Gly-Arg-CONH <sub>2</sub> BnSO <sub>2</sub> -D-(2,4-Dab)-Gly-Arg-CONH <sub>2</sub> BnSO <sub>2</sub> -D-Om-Gly-Arg-CONH <sub>2</sub>
25	BnSO <sub>2</sub> -D-homoLys-Gly-Arg-CONH <sub>2</sub> BnSO <sub>2</sub> -Bag-Gly-Arg-CONH <sub>2</sub> BnSO <sub>2</sub> -D-4-Gpa-Gly-Arg-CONH <sub>2</sub> BnSO <sub>2</sub> -D-3-Gpa-Gly-Arg-CONH <sub>2</sub> BnSO <sub>2</sub> -D-4-Apa-Gly-Arg-CONH <sub>2</sub>
30	BnSO <sub>2</sub> -D-3-Apa-Gly-Arg-CONH <sub>2</sub> BnSO <sub>2</sub> -D-4-Acg-Gly-Arg-CONH <sub>2</sub> BnSO <sub>2</sub> -D-(4-NH <sub>2</sub> Phe)-Gly-Arg-CONH <sub>2</sub> BnSO <sub>2</sub> -D-(3-NH <sub>2</sub> Phe)-Gly-Arg-CONH <sub>2</sub> BnSO <sub>2</sub> -D-Arg-Gly-(2,4-Dab)-H
35	BnSO <sub>2</sub> -D-Arg-Gly-(homoLys)-H BnSO <sub>2</sub> -D-Arg-Gly-(4-Gpa)-H

5	BnSO <sub>2</sub> -D-Arg-Gly-(3-Gpa)-H BnSO <sub>2</sub> -D-Arg-Gly-(4-Apa)-H BnSO <sub>2</sub> -D-Arg-Gly-(3-Apa)-H BnSO <sub>2</sub> -D-Arg-Gly-(4-Acg)-H BnSO <sub>2</sub> -D-Arg-Gly-(4-NH <sub>2</sub> Phe)-H
10	BnSO <sub>2</sub> -D-Arg-Gly-(3-NH <sub>2</sub> Phe)-H Boc-D-3-PIA-Gly-Arg-CONHEtPh Boc-D-4-PIA-Gly-Arg-CONHEtPh Boc-D-3-GPIA-Gly-Arg-CONHEtPh Boc-D-4-GPIA-Gly-Arg-CONHEtPh
15	Boc-D-3-PIG-Gly-Arg-CONHEtPh Boc-D-4-PIG-Gly-Arg-CONHEtPh Boc-D-3-GPIG-Gly-Arg-CONHEtPh Boc-D-4-GPIG-Gly-Arg-CONHEtPh BnSO <sub>2</sub> -D-Arg-Gly-Arg-CONHEtPh
20	BnSO <sub>2</sub> -D-3-PIA-Gly-Arg-CONHEtPh BnSO <sub>2</sub> -D-4-PIA-Gly-Arg-CONHEtPh BnSO <sub>2</sub> -D-3-GPIA-Gly-Arg-CONHEtPh BnSO <sub>2</sub> -D-4-GPIA-Gly-Arg-CONHEtPh BnSO <sub>2</sub> -D-3-PIG-Gly-Arg-CONHEtPh
25	BnSO <sub>2</sub> -D-4-PIG-Gly-Arg-CONHEtPh BnSO <sub>2</sub> -D-3-GPIG-Gly-Arg-CONHEtPh BnSO <sub>2</sub> -D-4-GPIG-Gly-Arg-CONHEtPh D-Arg-Gly-Arg-CONHEtPh D-3-PIA-Gly-Arg-CONHEtPh
30	D-4-PIA-Gly-Arg-CONHEtPh D-3-GPIA-Gly-Arg-CONHEtPh D-4-GPIA-Gly-Arg-CONHEtPh D-3-PIG-Gly-Arg-CONHEtPh D-4-PIG-Gly-Arg-CONHEtPh
35	D-3-GPIG-Gly-Arg-CONHEtPh D-4-GPIG-Gly-Arg-CONHEtPh BnSO <sub>2</sub> -D-Arg-Gly-Arg-CON(CH <sub>2</sub> ) <sub>4</sub> BnSO <sub>2</sub> -D-3-PIA-Gly-Arg-CON(CH <sub>2</sub> ) <sub>4</sub> BnSO <sub>2</sub> -D-4-PIA-Gly-Arg-CON(CH <sub>2</sub> ) <sub>4</sub>
40	BnSO <sub>2</sub> -D-3-GPIA-Gly-Arg-CON(CH <sub>2</sub> ) <sub>4</sub>



5	BnSO <sub>2</sub> -D-4-GPIA-Gly-Arg-CON(CH <sub>2</sub> ) <sub>4</sub> BnSO <sub>2</sub> -D-3-PIG-Gly-Arg-CON(CH <sub>2</sub> ) <sub>4</sub> BnSO <sub>2</sub> -D-4-PIG-Gly-Arg-CON(CH <sub>2</sub> ) <sub>4</sub> BnSO <sub>2</sub> -D-3-GPIG-Gly-Arg-CON(CH <sub>2</sub> ) <sub>4</sub> BnSO <sub>2</sub> -D-4-GPIG-Gly-Arg-CON(CH <sub>2</sub> ) <sub>4</sub>
10	Boc-D-Arg-Gly-Arg-CONHBn Boc-D-3-PIA-Gly-Arg-CONHBn Boc-D-4-PIA-Gly-Arg-CONHBn Boc-D-3-GPIA-Gly-Arg-CONHBn Boc-D-4-GPIA-Gly-Arg-CONHBn
15	Boc-D-3-PIG-Gly-Arg-CONHBn Boc-D-4-PIG-Gly-Arg-CONHBn Boc-D-3-GPIG-Gly-Arg-CONHBn Boc-D-4-GPIG-Gly-Arg-CONHBn Boc-D-Arg-Gly-Arg-CONHCH <sub>3</sub>
20	Boc-D-3-PIA-Gly-Arg-CONHCH <sub>3</sub> Boc-D-4-PIA-Gly-Arg-CONHCH <sub>3</sub> Boc-D-3-GPIA-Gly-Arg-CONHCH <sub>3</sub> Boc-D-4-GPIA-Gly-Arg-CONHCH <sub>3</sub> Boc-D-3-PIG-Gly-Arg-CONHCH <sub>3</sub>
25	Boc-D-4-PIG-Gly-Arg-CONHCH <sub>3</sub> Boc-D-3-GPIG-Gly-Arg-CONHCH <sub>3</sub> Boc-D-4-GPIG-Gly-Arg-CONHCH <sub>3</sub> D-Arg-Gly-Arg-CONHPh Boc-D-3-PIA-Gly-Arg-CONHPh
30	Boc-D-4-PIA-Gly-Arg-CONHPh Boc-D-3-GPIA-Gly-Arg-CONHPh Boc-D-4-GPIA-Gly-Arg-CONHPh Boc-D-3-PIG-Gly-Arg-CONHPh Boc-D-4-PIG-Gly-Arg-CONHPh
35	Boc-D-3-GPIG-Gly-Arg-CONHPh Boc-D-4-GPIG-Gly-Arg-CONHPh BnSO <sub>2</sub> -D-Arg-Gly-Arg-CONBn BnSO <sub>2</sub> -D-3-PIA-Gly-Arg-CONBn
40	

- 5                   BnSO<sub>2</sub>-D-4-PIA-Gly-Arg-CONBn  
                  BnSO<sub>2</sub>-D-3-GPIA-Gly-Arg-CONBn  
                  BnSO<sub>2</sub>-D-4-GPIA-Gly-Arg-CONBn  
                  BnSO<sub>2</sub>-D-3-PIG-Gly-Arg-CONBn  
                  BnSO<sub>2</sub>-D-4-PIG-Gly-Arg-CONBn  
10                  BnSO<sub>2</sub>-D-3-GPIG-Gly-Arg-CONBn  
                  BnSO<sub>2</sub>-D-4-GPIG-Gly-Arg-CONBn

The ketoamides synthesized in Examples 42-67, exhibited IC<sub>50</sub> values in the range of 4-79 nM for Factor Xa and from 5-165 nM for prothrombinase, with selectivities against thrombin of 10-1400.

- 15                  This invention also encompasses prodrug derivatives of the compounds contained herein. The term "prodrug" refers to a pharmacologically inactive derivative of a parent drug molecule that requires biotransformation, either spontaneous or enzymatic, within the organism to release the active drug. Prodrugs are variations or derivatives of the compounds of this invention which have  
20                  metabolically cleavable groups and become, by solvolysis under physiological conditions, or by enzymatic degradation the compounds of the invention which are pharmaceutically active *in vivo*. Prodrug compounds of this invention may be called  
25                  single, double, triple etc., depending on the number of biotransformation steps required to release the active drug within the organism, and indicating the number of  
                  functionalities present in a precursor-type form. Prodrug forms often offer advantages of solubility, tissue compatibility, or delayed release in the mammalian organism (see, Bundgard, H., Design of Prodrugs, pp. 7-9, 21-24, Elsevier, Amsterdam 1985 and Silverman, R.B., The Organic Chemistry of Drug Design and Drug Action, pp. 352-401, Academic Press, San Diego, CA, 1992). Prodrugs  
30                  commonly known in the art include acid derivatives well known to practitioners of the art, such as, for example, esters prepared by reaction of the parent acids with a suitable alcohol, or amides prepared by reaction of the parent acid compound with an amine, or basic groups reacted to form an acylated base derivative. Moreover, the prodrug derivatives of this invention may be combined with other features herein  
35                  taught to enhance bioavailability.

- In preferred embodiments of the present invention, basic groups such as guanidino or amidino functions are derivatized as carbamates or amides by acylation. Additionally, carbonyl functionalities are derivatized to form various acetals, enol esters, amins, thioacetals or Schiff's bases using alcohols, thiols,  
40                  amines or amides, carbamates, substituted hydrazines and the like.

## 5 Preparation of Compounds

The compounds of the present invention may be synthesized by either solid or liquid phase methods described and referenced in standard textbooks, or by a combination of both methods. These methods are well known in the art. See, Bodanszky, M., in "The Principles of Peptide Synthesis", Hafner, K., Rees, C.W.,  
10 Trost, B.M., Lehn, J.-M., Schleyer, P. v-R., Zahradnik, R., Eds., Springer-Verlag, Berlin, 1984. Starting materials are commercially available reagents and reactions are carried out in standard laboratory glassware and reaction vessels under reaction conditions of ambient temperature and pressure, except where otherwise indicated.

The peptide aldehyde compounds of the present invention may be  
15 synthesized by solution phase procedures described in United States Patent 5,380,713 of Balasubramanian *et al.*, or in the European Patent Application EPO 479 489 A2 of Gesellchen *et al.*, or by sequential chemical attachment of amino acid derivatives using the solid phase synthesis reagents and methods disclosed in WO 93/15756 or WO 94/13693.

20 Peptide alpha-keto acids, esters and amides can be prepared as described by Medhi, G. *et al.*, Biochem. Biophys. Res. Commun., 166, 595-600 (1990); Angelastro, *et al.*, J. Med. Chem., 33, 11-13 (1990); or by Webb, T.R. *et al.*, International Patent Application WO 9408941.

Peptide alpha-fluoroketones can be prepared as described by Neises, B.,  
25 European Patent Application EP 0504064 A1 published September 16, 1992 or described by Skiles, J.W. *et al.*, J. Med. Chem. 35, 641-662 (1992).


Peptide boronic acids and diesters can be prepared by the methods described by Kettner, C. *et al.*, J. Biol. Chem. 265, 18289-18297 (1990).

The starting materials used in any of these methods are commercially  
30 available from chemical vendors such as Aldrich, Sigma, Nova Biochemicals, Bachem Biosciences, and the like, or may be readily synthesized by known procedures.

During the synthesis of these compounds, the functional groups of the amino acid derivatives used in these methods are protected by blocking groups to prevent  
35 cross reaction during the coupling procedure. Examples of suitable blocking groups and their use are described in "The Peptides: Analysis, Synthesis, Biology", Academic Press, Vol. 3 (Gross, E. & Meienhofer, J., Eds., 1981) and Vol. 9 (1987), the disclosures of which are incorporated herein by reference.

- 5           Three exemplary synthesis schemes are outlined directly below, and the specific syntheses are described in the Examples. The reaction products are isolated and purified by conventional methods, typically by solvent extraction into a compatible solvent. The products may be further purified by column chromatography or other appropriate methods.





### Pharmaceutical and Diagnostic Compositions and Formulations

The compounds of this invention may be isolated as the free acid or base or converted to salts of various inorganic and organic acids and bases. Such salts are within the scope of this invention. Nontoxic and physiologically compatible salts are particularly useful although other less desirable salts may have use in the processes of isolation and purification.

A number of methods are useful for the preparation of the salts described above and are known to those skilled in the art. For example, reaction of the free acid or free base form of a compound of the structures recited above with one or more molar equivalents of the desired acid or base in a solvent or solvent mixture in which the salt is insoluble, or in a solvent like water after which the solvent is removed by evaporation, distillation or freeze drying. Alternatively, the free acid or base form of the product may be passed over an ion exchange resin to form the desired salt or one salt form of the product may be converted to another using the same general process.

5           Diagnostic applications of the compounds of this invention will typically utilize formulations such as solution or suspension. In the management of thrombotic disorders the compounds of this invention may be utilized in compositions such as tablets, capsules or elixirs for oral administration, suppositories, sterile solutions or suspensions or injectable administration, and the like, or incorporated into shaped  
10 articles. Subjects in need of treatment (typically mammalian) using the compounds of this invention can be administered dosages that will provide optimal efficacy. The dose and method of administration will vary from subject to subject and be dependent upon such factors as the type of mammal being treated, its sex, weight, diet, concurrent medication, overall clinical condition, the particular compounds  
15 employed, the specific use for which these compounds are employed, and other factors which those skilled in the medical arts will recognize.

Formulations of the compounds of this invention are prepared for storage or administration by mixing the compound having a desired degree of purity with physiologically acceptable carriers, excipients, stabilizers etc., and may be provided  
20 in sustained release or timed release formulations. Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical field, and are described, for example, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., (A.R. Gennaro edit. 1985). Such materials are nontoxic to the recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate,  
25 acetate and other organic acid salts, antioxidants such as ascorbic acid, low molecular weight (less than about ten residues) peptides such as polyarginine, proteins, such as serum albumin, gelatin, or immunoglobulins, hydrophilic polymers such as polyvinylpyrrolidinone, amino acids such as glycine, glutamic acid, aspartic acid, or arginine, monosaccharides, disaccharides, and other carbohydrates  
30 including cellulose or its derivatives, glucose, mannose or dextrans, chelating agents such as EDTA, sugar alcohols such as mannitol or sorbitol, counterions such as sodium and/or nonionic surfactants such as Tween, Pluronics or polyethyleneglycol.

Dosage formulations of the compounds of this invention to be used for therapeutic administration must be sterile. Sterility is readily accomplished by  
35 filtration through sterile membranes such as 0.2 micron membranes, or by other conventional methods. Formulations typically will be stored in lyophilized form or as an aqueous solution. The pH of the preparations of this invention typically will be between 3 and 11, more preferably from 5 to 9 and most preferably from 7 to 8. While the preferred route of administration is by injection such as intravenously  
40 (bolus and/or infusion), other methods of administration are also anticipated,

5 subcutaneously, intramuscularly, colonically, rectally, nasally or intraperitoneally, employing a variety of dosage forms such as suppositories, implanted pellets or small cylinders, aerosols, oral dosage formulations and topical formulations such as ointments, drops and dermal patches. The compounds of this invention are desirably incorporated into shaped articles such as implants which may employ inert  
10 materials such as biodegradable polymers or synthetic silicones, for example, Silastic, silicone rubber or other polymers commercially available.

The compounds of this invention may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of  
15 lipids, such as cholesterol, stearylamine or phosphatidylcholines.

The compounds of this invention may also be delivered by the use of antibodies, antibody fragments, growth factors, hormones, or other targeting moieties, to which the compound molecules are coupled. The compounds of this invention may also be coupled with suitable polymers as targetable drug carriers.  
20 Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropyl-methacrylamide-phenol, polyhydroxyethyl-aspartamide-phenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the factor Xa inhibitors of this invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example polylactic  
25 acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross linked or amphipathic block copolymers of hydrogels. Polymers and semipermeable polymer matrices may be formed into shaped articles, such as valves, stents, tubing, prostheses and the like.  
30 Therapeutic compound liquid formulations generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by hypodermic injection needle.

Therapeutically effective dosages may be determined by either *in vitro* or *in vivo* methods. For each particular compound of the present invention, individual  
35 determinations may be made to determine the optimal dosage required. The range of therapeutically effective dosages will naturally be influenced by the route of administration, the therapeutic objectives, and the condition of the patient. For injection by hypodermic needle, it may be assumed the dosage is delivered into the body's fluids. For other routes of administration, the absorption efficiency must be  
40 individually determined for each inhibitor by methods well known in pharmacology. Accordingly, it may be necessary for the therapist to titer the dosage and modify the



5 route of administration as required to obtain the optimal therapeutic effect. The determination of effective dosage levels, that is, the dosage levels necessary to achieve the desired result, will be within the ambit of one skilled in the art. Typically, applications of compound are commenced at lower dosage levels, with dosage levels being increased until the desired effect is achieved.

10 A typical dosage might range from about 0.001 mg/kg to about 1000 mg/kg, preferably from about 0.01 mg/kg to about 100 mg/kg, and more preferably from about 0.10 mg/kg to about 20 mg/kg. Advantageously, the compounds of this invention may be administered several times daily, and other dosage regimens may also be useful.

15 Typically, about 0.5 to 500 mg of a compound or mixture of compounds of this invention, as the free acid or base form or as a pharmaceutically acceptable salt, is compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, dye, flavor etc., as called for by accepted pharmaceutical practice. The amount of active ingredient in these compositions is  
20 such that a suitable dosage in the range indicated is obtained.

Typical adjuvants which may be incorporated into tablets, capsules and the like are a binder such as acacia, corn starch or gelatin, and excipient such as microcrystalline cellulose, a disintegrating agent like corn starch or alginic acid, a lubricant such as magnesium stearate, a sweetening agent such as sucrose or  
25 lactose, or a flavoring agent. When a dosage form is a capsule, in addition to the above materials it may also contain a liquid carrier such as water, saline, a fatty oil. Other materials of various types may be used as coatings or as modifiers of the physical form of the dosage unit. Sterile compositions for injection can be formulated according to conventional pharmaceutical practice. For example,  
30 dissolution or suspension of the active compound in a vehicle such as an oil or a synthetic fatty vehicle like ethyl oleate, or into a liposome may be desired. Buffers, preservatives, antioxidants and the like can be incorporated according to accepted pharmaceutical practice.

In practicing the methods of this invention, the compounds of this invention  
35 may be used alone or in combination, or in combination with other therapeutic or diagnostic agents. In certain preferred embodiments, the compounds of this invention may be coadministered along with other compounds typically prescribed for these conditions according to generally accepted medical practice, such as anticoagulant agents, thrombolytic agents, or other antithrombotics, including  
40 platelet aggregation inhibitors, tissue plasminogen activators, urokinase, prourokinase, streptokinase, heparin, aspirin, or warfarin. The compounds of this

- 5 invention can be utilized *in vivo*, ordinarily in mammals, such as humans, sheep, horses, cattle, pigs, dogs, cats, rats and mice, or *in vitro*.

The preferred compounds of the present invention are characterized by their ability to inhibit thrombus formation with acceptable effects on classical measures of coagulation parameters, platelets and platelet function, and acceptable levels of  
10 bleeding complications associated with their use. Conditions characterized by undesired thrombosis would include those involving the arterial and venous vasculature.

With respect to the coronary arterial vasculature, abnormal thrombus formation characterizes the rupture of an established atherosclerotic plaque which  
15 is the major cause of acute myocardial infarction and unstable angina, as well as also characterizing the occlusive coronary thrombus formation resulting from either thrombolytic therapy or percutaneous transluminal coronary angioplasty (PTCA).

With respect to the venous vasculature, abnormal thrombus formation characterizes the condition observed in patients undergoing major surgery in the  
20 lower extremities or the abdominal area who often suffer from thrombus formation in the venous vasculature resulting in reduced blood flow to the affected extremity and a predisposition to pulmonary embolism. Abnormal thrombus formation further characterizes disseminated intravascular coagulopathy which commonly occurs within both vascular systems during septic shock, certain viral infections and cancer, and is  
25 a condition where there is rapid consumption of coagulation factors and systemic coagulation which results in the formation of life-threatening thrombi occurring throughout the microvasculature leading to widespread organ failure.

The compounds of this present invention, selected and used as disclosed herein, are believed to be useful for preventing or treating a condition characterized  
30 by undesired thrombosis, such as in (a) the treatment or prevention of any thrombotically mediated acute coronary syndrome including myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, (b) the treatment or prevention of any thrombotically mediated cerebrovascular syndrome including embolic stroke,  
35 thrombotic stroke or transient ischemic attacks, (c) the treatment or prevention of any thrombotic syndrome occurring in the venous system including deep venous thrombosis or pulmonary embolus occurring either spontaneously or in the setting of malignancy, surgery or trauma, (d) the treatment or prevention of any coagulopathy including disseminated intravascular coagulation (including the setting of septic  
40 shock or other infection, surgery, pregnancy, trauma or malignancy and whether associated with multi-organ failure or not), thrombotic thrombocytopenic purpura,

- 5 thromboangiitis obliterans, or thrombotic disease associated with heparin induced thrombocytopenia, (e) the treatment or prevention of thrombotic complications associated with extracorporeal circulation (e.g. renal dialysis, cardiopulmonary bypass or other oxygenation procedure, plasmapheresis), (f) the treatment or prevention of thrombotic complications associated with instrumentation (e.g. cardiac
- 10 or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve), and (g) those involved with the fitting of prosthetic devices.

Anticoagulant therapy is also useful to prevent coagulation of stored whole blood and to prevent coagulation in other biological samples for testing or storage. Thus the compounds of this invention can be added to or contacted with any

15 medium containing or suspected to contain factor Xa and in which it is desired that blood coagulation be inhibited, e.g., when contacting the mammal's blood with material such as vascular grafts, stents, orthopedic prostheses, cardiac stents, valves and prostheses, extra corporeal circulation systems and the like.


Without further description, it is believed that one of ordinary skill in the art

20 can, using the preceding description and the following illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods. The following working examples therefore, specifically point out preferred embodiments of the present invention, and are not to be construed as limiting in any way the remainder of the disclosure.

25

### EXAMPLES

#### EXAMPLE 1



#### 30 Boc-D-Lys(Cbz)-Gly-OH

To a solution of 1.14 g (3.0 mmol) of Boc-D-Lys(Cbz)-OH and 0.34 g (3.0 mmol) of N-hydroxysuccinimide dissolved in 20 ml of dioxane was added a solution of 0.62 g (3.0 mmol) of DCC dissolved in 4.0 ml of dioxane. The resultant solution was stirred at room temperature under a nitrogen atmosphere for 18 hrs, during

35 which time the DCU separated from the solution. After removal of the solids by vacuum filtration, the clear colorless filtrate was treated with a single portion of glycine (0.36 g, 4.9 mmol) and sodium bicarbonate (0.90 g, 10.8 mmol) dissolved in 30 ml of water. The resultant mixture was stirred at room temperature for 2 days.


- 5 The solution was then acidified to pH=1-2 with 10% HCl solution and extracted with ethyl acetate three times. The combined organic extracts were washed twice with brine, dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to afford 1.30 g (3.0 mmol, 100% yield) of the product as a colorless, viscous oil.

RP-HPLC: retention 13.12 min; C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0  
10 ml/min.

NMR (CDCl<sub>3</sub>): 7.22 (s, C<sub>6</sub>H<sub>5</sub>), 4.98 (s, OCH<sub>2</sub>Ar), 3.95-4.15 (m, NHCH<sub>2</sub>COOH), 3.80-3.93 (m, NHCHCO), 2.94-3.10 (m, CH<sub>2</sub>NHCO), 1.00-1.60 (m, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.30 (s, (CH<sub>3</sub>)<sub>3</sub>CO).

15

### EXAMPLE 2



#### Boc-Arg(Cbz)-lactam

- A solution of Boc-Arg(Cbz)-OH (10.6 g, 26.0 mmol) dissolved in 70 ml of dry THF was cooled in a dry ice-acetone bath under a N<sub>2</sub> atmosphere. N-Methylmorpholine (6.5 ml, 59.1 mmol) was added to the solution followed by isobutyl chloroformate (3.8 ml, 29.3 mmol). A white solid separated immediately, and the suspension was stirred at the temperature of the dry ice-acetone bath for 1 hr, warmed to room temperature and stirred an additional 2.5 hrs. The mixture was poured onto ice-water and the precipitated solid collected and dried *in vacuo*. There  
20 was obtained 8.35 g (21.4 mmol, 82% yield) of the product as a white solid.


RP-HPLC: retention 15.09 min; C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0  
ml/min.

- NMR (CDCl<sub>3</sub>): 9.4-9.7 (bm, NH), 7.30-7.50 (m, C<sub>6</sub>H<sub>5</sub>), 5.25-5.35 (d, NHCH), 5.20 (s, OCH<sub>2</sub>Ar), 4.80-5.00 (m, NCHCH<sub>2</sub>), 3.30-3.60/4.30-4.60 (m, NCH<sub>2</sub>), 1.80-2.10/2.40-2.60 (m, CH<sub>2</sub>CH<sub>2</sub>), 1.50 (s, (CH<sub>3</sub>)<sub>3</sub>CO).

30

IR: 3346 cm<sup>-1</sup>, 3273 cm<sup>-1</sup>, 3244 cm<sup>-1</sup>, 1726 cm<sup>-1</sup>, 1698 cm<sup>-1</sup>, 1644 cm<sup>-1</sup>.

### EXAMPLE 3




35

5 Arg(Cbz)-lactam·HCl salt

Boc-Arg(Cbz)-lactam (1.20g, 3.1 mmol) prepared in Example 2 was treated with 10 ml of 4N HCl/dioxane for 1 hr at room temperature under N<sub>2</sub>. Removal of solvent *in vacuo* produced a white glass. This material was triturated with ether and filtered to afford 1.05 g (3.2 mmol, 100% yield) of a non-hygroscopic white powder.

10 RP HPLC: retention time 11.34 min; C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 ml/min.

EXAMPLE 415 Boc-D-Lys(Cbz)-Gly-Arg(Cbz)-lactam

To a solution of 1.30 g (3.0 mmol) of Boc-D-Lys(Cbz)-Gly-OH prepared in Example 1, in 30 ml of dioxane was added N-hydroxysuccinimide (0.35 g, 3.0 mmol), followed by a solution of DCC (0.63 g, 3.0 mmol) in 5 ml of dioxane. The resultant solution was stirred at room temperature under N<sub>2</sub> for 17 hrs, during which

20 time DCU precipitated. After removal of solids by vacuum filtration, the clear, colorless filtrate was concentrated to dryness *in vacuo* to yield a colorless oil. This material was dissolved in 25 ml of DMF and treated with the Arg(Cbz)-lactam·HCl salt of Example 3 (0.98 g, 3.0 mmol) which was added as a solid followed by the addition of 25 ml of DMF. After the lactam had dissolved completely, N-

25 methylmorpholine (1.0 ml, 9.1 mmol) was added to neutralize the lactam salt. The clear colorless solution was stirred at room temperature under N<sub>2</sub> for 25 hrs. The resultant yellow solution was acidified to pH=1-2 with 10% HCl solution and extracted twice with ethyl acetate. The combined organic extracts were washed six times with brine, dried over MgSO<sub>4</sub> and concentrated *in vacuo*. There was obtained


30 1.39 g (2.0 mmol, 65% yield) of a white solid product.

RP-HPLC: retention 14.46 min; C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.5 ml/min.

35 MS: 710 by thermospray; calculated (M+H) = 710.8.

EXAMPLE 5

SUBSTITUTE SHEET (RULE 26)



5

Boc-D-Lys(Cbz)-Gly-Arg(Cbz)-H


To a dry flask equipped with a N<sub>2</sub> inlet, low temperature thermometer, and rubber septum was introduced a solution of Boc-D-Lys(Cbz)-Gly-Arg(Cbz)-lactam (1.37 g, 1.9 mmol) prepared in Example 4, dissolved in 50 ml of anhydrous THF.

- 10 The flask was cooled in a CCl<sub>4</sub>-dry ice bath such that the internal temperature was maintained between -35°C and -25°C. To the cooled solution was added LiAlH<sub>4</sub>/THF (1.0 M solution) (1.9 ml, 1.9 mmol). The resultant solution was stirred in the cold under N<sub>2</sub> for 2 hr. After this time, the reaction was quenched by introduction of 5 ml of CH<sub>3</sub>OH to the chilled solution, followed by gradual warming
- 15 to ambient temperature. Sulfuric acid solution (1 N, 36 ml) was added and extracted with ethyl acetate. The organic phase was washed twice with brine, dried over MgSO<sub>4</sub>, and concentrated *in vacuo*. There was obtained 1.23 g (1.7 mmol, 89% yield) of the product as a white solid.

RP-HPLC: retention 13.34 min; retention time 13.41 min

- 20 (thiosemicarbazone); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 ml/min.

MS: M+H=712 by thermospray ; calculated (M+H) = 712.8.

EXAMPLE 6

- 25 Boc-D-Lys-Gly-Arg-H

- A Parr bottle was charged with Boc-D-Lys(Cbz)-Gly-Arg(Cbz)-H prepared in Example 5, (1.21 g, 1.7 mmol), 10% palladium on carbon (0.43 g, 0.4 mmol) and 35 ml of absolute ethanol. Hydrogenation was conducted at 40 psi in a Parr shaker for
- 30 24 hrs after which the catalyst was removed by vacuum filtration over Celite. The clear colorless filtrate was diluted with water and acetic acid (1 ml) and lyophilized. The lyophilized powder was purified by preparative RP-HPLC using a gradient from 10% CH<sub>3</sub>CN in water (containing 0.1% TFA) to 50% CH<sub>3</sub>CN in water (containing


- 5 0.1% TFA) using a flow rate of 10 ml/min. The desired fractions were combined and lyophilized to yield 0.16 g (0.35 mmol, 20% yield) of the product as a white powder.

RP-HPLC: retention 8.66 min and 9.26 min (tautomers); retention time 8,7 min (thiosemicarbazone); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 ml/min.

MS: M+H=444 by thermospray; calculated (M+H) = 444.5.

10

#### EXAMPLE 7



#### Cbz-D-Arg(Cbz)<sub>2</sub>-Gly-OH


- Cbz-D-Arg(Cbz)<sub>2</sub>-Gly-OH is prepared according to Example 1 by coupling  
15 the N-hydroxysuccinimide active ester of Cbz-D-Arg(Cbz)<sub>2</sub>-OH to glycine in a yield of 87%.

RP-HPLC: retention 14.09 min; C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 ml/min.

MS: (M+H)= 634.4 by electrospray; calculated (M+H)= 634.7.

20

#### EXAMPLE 8



#### BnOSuc-D-Arg(Cbz)<sub>2</sub>-OH

- To a solution of 186 mg (0.89 mmol) of succinic acid monobenzyloxycarbonyl ester and  
25 113 mg (1.0 mmol) of N-hydroxysuccinimide in 1.7 mL of p-dioxane is added 194 mg (0.93 mmol) of DCC in 0.5 mL of dioxane. The solution is allowed to stir for 18hr. The white precipitate of DCU is filtered and the solution is added dropwise to 4.5 mL of 0.6M NaHCO<sub>3</sub> containing 485 mg (0.89 mmol) of H-D-Arg(Cbz)<sub>2</sub>-OH (prepared by treatment of Boc-D-Arg(Cbz)<sub>2</sub>-OH with TFA for 1h followed by  
30 concentration *in vacuo*). After stirring for 24h at 23°C, the reaction was diluted with 1N HCl and extracted 3 times with EtOAc. The organic layers were combined and


- 5 washed with brine, dried (MgSO<sub>4</sub>), and concentrated (in vacuo) to give 516 mg (91%) of a clear oil.

NMR-90MHz (CDCl<sub>3</sub>): 7.6-7.2 (br s, 15), 5.3 (s, 2), 5.2 (s, 2), 5.1 (s, 2), 4.5 (m, 1), 3.5 (br m, 2), 2.4 (br m, 4), 1.8 (br m, 4).

- RP-HPLC: retention 14.94 min; C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 ml/min.

MS: (M+H)= 633.6 by electrospray; calculated (M+H)= 633.7.

#### EXAMPLE 9




#### 15 BnOSuc-(D-Arg)<sub>2</sub>-Gly-OH

O-benzylsuccinoyl-D-arginyl(Cbz<sub>2</sub>)-glycine is prepared according to Example 1 by coupling the N-hydroxysuccinimide active ester of O-benzylsuccinoyl-D-arginine(Cbz<sub>2</sub>)-OH to glycine in a crude yield of 81%. This material was purified by flash silica gel chromatography eluting with 0, 2.5, 5, 7.5, and 10% methanol/DCM.

- 20 RP-HPLC: retention 13.92 min; C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 ml/min.

MS: (M+H)= 690.8 by electrospray; calculated (M+H)= 690.7.

#### EXAMPLE 10



#### 25 Boc-Gly-Arg(N-Cbz)-lactam


- To a solution of Boc-Gly (173 mg, 1.1 mmol) and the Arg(Cbz)-lactam·HCl salt of Example 3 (330 mg, 1.0 mmol) in 2 mL of 1:1 hexane/THF at 0°C is added BOP (472 mg, 1.0mmol) followed by 191 uL of DIEA (1.1 mmol). The solution is allowed to warm to 23°C and stir for 3 hrs at which time the solvent is removed *in vacuo*. The residue is dissolved in 10 mL of EtOAc and extracted twice with sat. NaHCO<sub>3</sub>, twice with 5% H<sub>3</sub>PO<sub>4</sub>, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated (*in vacuo*) to afford a 80-95% yield of the product as a white solid.



5 RP-HPLC retention, 13.92 min, linear gradient: 30% water/TEAP(pH 2.5) to 100% CH<sub>3</sub>CN over 30 min at 1.5 mL/min.

MS: (M+H)= 873.0 by electrospray; calculated (M+H)= 873.0.

### EXAMPLE 11



10

#### Boc-D-Arg-(Cbz<sub>2</sub>)-Gly-Arg(N-Cbz)-lactam

Boc-D-Arg(Cbz<sub>2</sub>)-Gly-Arg(N-Cbz)-lactam is prepared by coupling H-Gly-Arg(Cbz<sub>2</sub>)-lactam (prepared by deprotection of Boc-Gly-Arg(Cbz<sub>2</sub>)-lactam with 40% TFA/DCM at 23°C over 1h and concentration *in vacuo*) to the tert-butylloxycarbonyl-D-Arg(Cbz<sub>2</sub>)-OH according to the procedure in example 10. The product was obtained in 90% yield as a white solid.


15

RP-HPLC retention, 13.92 min, linear gradient: 30% H<sub>2</sub>O/TEAP(1% TEA/1% H<sub>3</sub>PO<sub>4</sub>, pH 2.5) to 100% CH<sub>3</sub>CN over 30 min at 1.5 mL/min.

MS: (M+H)= 873.0 by electrospray; calculated (M+H)= 873.0.

20

### EXAMPLE 12



#### Cbz-D-Arg(Cbz<sub>2</sub>)-Gly-Arg(Cbz)-lactam


25 Benzyloxycarbonyl-D-Arg(Cbz<sub>2</sub>)-Gly-Arg(Cbz)-lactam is prepared by coupling H-Arg(Cbz<sub>2</sub>)-lactam to the N-hydroxysuccinimide active ester of benzyloxycarbonyl-D-Arg(Cbz<sub>2</sub>)-Gly-OH according to the procedure in Example 4. The crude oil obtained after extraction was purified by flash silica gel chromatography eluting with 0, 2.5, 5, 6% MeOH/DCM. The appropriate fractions were combined and concentrated *in vacuo* to yield 37% of a white solid:

30

RP-HPLC retention, 14.60 min; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

5 MS: (M+H)= 906.6 by electrospray; calculated (M+H)= 906.9.

### EXAMPLE 13



#### EtO<sub>2</sub>C-D-Arg(Cbz<sub>2</sub>)-Gly-Arg(Cbz)-lactam

10 Ethoxycarbonyl-D-Arg(Cbz<sub>2</sub>)-Gly-Arg-(Cbz)-lactam is prepared by coupling H-Arg(Cbz<sub>2</sub>)-lactam to the N-hydroxysuccinimide active ester of ethoxycarbonyl-D-Arg-Gly-OH according to the procedure in Example 4. The crude oil obtained after extraction was purified by flash silica gel chromatography eluting with 0, 2.5, 5, 10% MeOH/DCM. The appropriate fractions were combined and concentrated *in vacuo*


15 to yield 46% of a white solid:

RP-HPLC retention, 14.13 min; C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

MS: (M+H)= 844.1 and (M+H+thioglycerol)= 952.1 by fast atom bombardment; calculated (M+H)= 844.4.

20

### EXAMPLE 14



#### O-Benzylsuccinoyl-D-Arg(Cbz<sub>2</sub>)-Gly-Arg(Cbz)-lactam


25 O-Benzylsuccinoyl-D-Arg(Cbz<sub>2</sub>)-Gly-Arg(Cbz)-lactam is prepared by coupling of H-Arg(Cbz<sub>2</sub>)-lactam to the succinimide active ester of O-benzylsuccinoyl-D-Arg(Cbz<sub>2</sub>)-Gly-OH according to the procedure in Example 4. The crude oil obtained after extraction was purified by flash silica gel chromatography eluting with 0, 2.5, 5, 10% MeOH/DCM. The appropriate fractions were combined and

30 concentrated *in vacuo* to yield 41% of a white solid:

RP-HPLC retention, 14.54 min; C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

5 MS: (M+H)= 963.1 by electrospray; calculated (M+H)= 963.06.

### EXAMPLE 15



#### Cbz-D-Arg(Cbz<sub>2</sub>)-Gly-Arg(Cbz)-H


10 Cbz-D-Arg(Cbz<sub>2</sub>)-Gly-Arg(Cbz)-H is prepared by reduction of the corresponding lactam in 77% yield according to the procedure in Example 5.

RP-HPLC: retention 13.9 and 14.1 min (tautomers); 14.2 min (thiosemicarbazone derivative); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

MS: (M+H)= 908.6 and (M+NH<sub>4</sub>)= 926.8, calculated (M+H)= 908.9.

15

### EXAMPLE 16



#### EtO<sub>2</sub>C-D-Arg(Cbz<sub>2</sub>)-Gly-Arg(Cbz)-H

20 EtO<sub>2</sub>C-D-Arg(Cbz<sub>2</sub>)-Gly-Arg(Cbz)-H is prepared by reduction of the corresponding lactam as prepared in Example 13 in 68% yield according to the procedure in Example 5.


RP-HPLC retention, 13.9 and 14.1 min (tautomers); 14.2 min (thiosemicarbazone derivative); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

25

MS: (M+H)= 846.5 and (M+NH<sub>4</sub>)= 864.5 by fast atom bombardment, calculated (M+H)= 846.9.

### EXAMPLE 17

42



5

O-Benzylsuccinoyl-D-Arg(Cbz<sub>2</sub>)-Gly-Arg(Cbz)-H


O-Benzylsuccinoyl-D-Arg(Cbz<sub>2</sub>)-Gly-Arg(Cbz)-H is prepared by reduction of the corresponding lactam, as prepared in Example 14 in 89% yield according to the procedure in Example 5.

10

RP-HPLC retention, 13.74 and 13.97 min (tautomers), 13.87 min (thiosemicarbazone derivative); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

MS: (M+H)= 965.2 by electrospray, calculated (M+H)= 965.1.

EXAMPLE 18



15 Boc-D-Arg-Gly-Arg-H

Boc-D-Arg-Gly-Arg-H was prepared by hydrogenolysis of the corresponding protected peptide aldehyde Cbz-D-Arg(Cbz<sub>2</sub>)-Gly-Arg(Cbz)-H (prepared by reduction of the corresponding lactam of Example 11 in 77% yield according to the procedure in Example 5 and deprotected directly without further purification or characterization) following the procedure in Example 6 except that THF/hexane was used as solvent. After 1h, the deprotection is complete and the catalyst is removed by vacuum filtration through Celite. The compound is then purified by preparative reversed-phase HPLC using a gradient of 100% water/0.1% TFA to 50% CH<sub>3</sub>CN/0.1% TFA. The appropriate fractions are pooled and lyophilized to give the desired product as white solid.


25

RP-HPLC retention: 8.58 and 8.65 min (tautomers); 8.66 min (semicarbazone derivative); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

MS: (M+H)= 472.3 by fast atom bombardment, calculated (M+H)= 472.3.

30

EXAMPLE 19




5

D-Arg-Gly-Arg-H

D-Arg-Gly-Arg-H was prepared by hydrogenolysis of the corresponding protected peptide aldehyde Cbz-D-Arg(Cbz<sub>2</sub>)-Gly-Arg(Cbz)-H as prepared in Example 15 following the procedure in Example 6. After 1 h, the deprotection is complete and the catalyst is removed by vacuum filtration through Celite. The pH of the solution is adjusted to 5 using Biorad WGR-2 (free amine form), filtered and lyophilized to give the desired product as white solid in 84 % yield.

RP-HPLC retention: 2.55 min ; 2.87 min (thiosemicarbazone derivative); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

MS: (M+H)= 372.1 by electrospray, calculated (M+H)= 372.5.

EXAMPLE 20EtO<sub>2</sub>C-D-Arg-Gly-Arg-H


EtO<sub>2</sub>C-D-Arg-Gly-Arg-H was prepared by hydrogenolysis of the corresponding protected peptide aldehyde of Example 16 following the procedure in Example 6 except that 0.1M THF and 0.2M aqueous acetic acid was used as solvent. After 1 h, the deprotection is complete and the catalyst is removed by vacuum filtration through Celite. The solution is lyophilized to give the desired product as white solid in 71 % yield.

RP-HPLC retention, 8.5 and 8.7 min (tautomers) ; 8.9 min (thiosemicarbazone derivative); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

MS: (M+H)= 444.2 by fast atom bombardment, calculated (M+H)= 444.3.

30

EXAMPLE 21



5

Succinoyl-D-Arg-Gly-Arg-H


Succinoyl-D-Arg-Gly-Arg-H was prepared by hydrogenolysis of the corresponding protected peptide aldehyde of Example 17 following the procedure in Example 6. After 1 h, the deprotection is complete and the catalyst is removed by  
10 filtration through Celite. The solution is lyophilized to give the desired product as white solid in 49 % yield.

RP-HPLC retention: 3.02 min ;3.51 min (thiosemicarbazone derivative);  
12.55 min (2,4 dinitrophenyl-hydrazone derivative); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25  
minutes, 2.0 mL/min.

15

MS: (M+H)= 472.8 by electrospray ,calculated (M+H)= 472.5.


5

EXAMPLE 22Benzyl trans-4-aminomethylcyclohexane-1-carboxylate p-toluenesulfonic acid salt

A mixture of 10.0 g (63.6 mmol) of trans-4-aminomethylcyclohexane-1-carboxylic acid, 25.1 g (132 mmol) of 4-toluenesulfonic acid, 50 mL (482 mmol) of benzyl alcohol, and 75 mL of toluene was heated to reflux under a N<sub>2</sub> atmosphere using a Dean-Stark trap to collect water formed during the reaction. During reflux, all solids dissolved to leave a colorless solution. After 3 hours at reflux, the solution began to turn yellow. Heat was removed, and the flask allowed to cool to room temperature. The solids which separated out were collected and washed with diethyl ether. There was obtained 24.5 g (58.4 mmol, 92% yield) of product as a white solid.

NMR (CD<sub>3</sub>OD): 7.12 / 7.20 / 7.65 / 7.73 (ABq, C<sub>6</sub>H<sub>4</sub>); 7.26 (s, C<sub>6</sub>H<sub>5</sub>); 5.06 (s, OCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>); 2.72 / 2.80 (d, CH<sub>2</sub>N); 2.32 (s, ArCH<sub>3</sub>); 1.00-2.18 (m, cyclohexyl H)

MS: M<sup>+</sup> = 247 by electron-impact; calculated M<sup>+</sup> = 247.3.

EXAMPLE 23Benzyl trans-4-(tert-butoxycarbonylhydrazidocarbonyl)-aminomethylcyclohexane-1-carboxylate:

To a solution of 3.24 g (20.0 mmol) of carbonyl diimidazole dissolved in 50 mL of DMF was added over 40 minutes a solution of 2.65 g (20.0 mmol) of tert-butylcarbazate dissolved in 50 mL of DMF. The resultant solution was stirred at room temperature under a N<sub>2</sub> atmosphere for 10 minutes. A solution of 7.16 g (17.1 mmol) of benzyl trans-4-aminomethylcyclohexane-1-carboxylate-TsOH dissolved in 50 mL of DMF was then added over 65 minutes. Finally, after the above solution was stirred for 5 min, 10 mL (71.7 mmol) of triethylamine was added over a 5 minute period. The resultant light yellow solution was stirred for 18 1/2 hrs.


5           The solution was acidified with the addition of 10% HCl, transferred to a separatory funnel, and extracted twice with ethyl acetate. The combined organic extracts were washed three times with saturated brine, dried over MgSO<sub>4</sub>, and concentrated *in vacuo*. Under high vacuum, there was obtained 6.64 g (16.4 mmol, 96% yield) of product as a sticky, yellow foam.

10           RP-HPLC: retention time = 13.73 min; C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

          NMR (CDCl<sub>3</sub>): 7.25 (s, C<sub>6</sub>H<sub>5</sub>); 6.67 / 6.72 (2 s, NH x 2); 5.50-5.62 (t, NHCH<sub>2</sub>); 5.00 (s, OCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>); 2.95-3.05 (t, CH<sub>2</sub>N); 0.80-2.20 (m, cyclohexyl H); 1.36 (s, (CH<sub>3</sub>)<sub>3</sub>CO)

15

#### EXAMPLE 24



trans-4-(tert-butoxycarbonylhydrazidocarbonyl)aminomethylcyclohexane-1-carboxylic acid


20           A Parr Bottle was charged with 5.46 g (13.2 mmol) of benzyl trans-4-(tert-butoxycarbonylhydrazidocarbonyl)-aminomethylcyclohexane-1-carboxylate dissolved in 60 mL of methanol. Added was 0.50 g (0.47 mmol) of 10% palladium on carbon catalyst. After hydrogenation at 38 psi on a Parr shaker for 90 minutes, the catalyst was removed by vacuum filtration over Celite. Concentration *in vacuo*  
25   of the filtrate gave 4.22 g (13.3 mmol, 100%) of product as a white solid.

IR: 3308 cm<sup>-1</sup>, 3276 cm<sup>-1</sup>, 1732 cm<sup>-1</sup>, 1720 cm<sup>-1</sup>, 1704 cm<sup>-1</sup>

MS: M+H = 316 by DCI; calculated (M+H) = 316.3.



5

EXAMPLE 25


trans-4-(hydrazidocarbonyl)aminomethylcyclohexane-1-carboxylic acid,  
trifluoroacetate salt:

A solution of 3.92 g (12.4 mmol) of trans-4-  
 10 (tert-butoxycarbonylhydrazidocarbonyl)-aminomethylcyclohexane-1-carboxylic acid  
 dissolved in 25 mL of TFA and 25 mL of DCM was stirred at room temperature  
 under a N<sub>2</sub> atmosphere for 90 minutes. Solvent was removed *in vacuo* to leave a  
 viscous, pale yellow oil. Addition of ethanol induced solidification. The solid was  
 collected and washed with diethyl ether to produce 2.19 g (6.65 mmol, 54% yield) of  
 15 product as a white solid.

RP-HPLC: retention time = 2.87 minutes; C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25  
 minutes, 2.0 mL/min.

IR: 3357 cm<sup>-1</sup>, 3264 cm<sup>-1</sup>, 1705 cm<sup>-1</sup>, 1666 cm<sup>-1</sup>

20

EXAMPLE 26

Boc-Arg(Cbz)<sub>2</sub>-H:

A solution of 2.72 g (5.00 mmol) of Boc-Arg(Cbz)<sub>2</sub>-OH and 0.81 g (5.00  
 mmol) of CDI dissolved in 40 mL of anhydrous THF was stirred at room temperature  
 25 under a N<sub>2</sub> atmosphere for 3 hrs. The flask was cooled in a dry ice/acetone bath  
 and 5.0 mL (5.00 mmol) of 1 Molar LiAlH<sub>4</sub> (in THF) was added by syringe. The  
 solution was stirred in the cold bath for 3 hours, then quenched by addition of 10  
 mL of 10% HCl solution. The flask was then removed from the cold bath.

After the flask warmed to room temperature, the contents were transferred to  
 30 a separatory funnel, diluted with additional 10% HCl and extracted twice with ethyl  
 acetate. The combined organic extracts were washed three times with saturated


- 5 brine, dried over  $\text{MgSO}_4$ , and concentrated *in vacuo*. Under high vacuum, there was obtained 2.69 g (5.10 mmol, 102% yield) of product as a white foam.

RP-HPLC: retention time = 13.28 min; 13.43 min (thiosemicarbazone);  
C<sub>18</sub>, O-100%  $\text{CH}_3\text{CN}$  over 25 minutes, 2.0 mL/min.

MS:  $\text{M}+\text{H} = 527$  by DCI; calculated  $(\text{M}+\text{H}) = 527.5$ .

10

#### EXAMPLE 27




N $\alpha$ -Boc(Cbz<sub>2</sub>)Argininal semicarbazonyl-N<sub>2</sub>-trans-4-methylenecyclohexane-1-carboxylic acid:

- 15 A solution of 2.69 g (5.10 mmol) of Boc-Arg(Cbz)<sub>2</sub>-H, 2.17 g (6.60 mmol) of trans-4-(hydrazidocarbonyl)aminomethylcyclohexane-1-carboxylic acid trifluoroacetate salt, 1.30 g (9.53 mmol) of sodium acetate, 75 mL of ethanol, and 75 mL of water was refluxed for 2 1/4 hrs. The solution was allowed to cool to room temperature, during which time the flask contents became cloudy.
- 20 After acidification to pH=1 with 50 mL of 10% HCl, the solution was extracted three times with ethyl acetate. The combined organic extracts were washed once with saturated brine solution, dried over  $\text{MgSO}_4$ , and concentrated *in vacuo*. Under high vacuum, there was obtained 2.78 g (3.84 mmol, 75% yield) of product as a pale yellow, microcrystalline foam.
- 25 RP-HPLC: retention time = 14.11 minutes; C<sub>18</sub>, O-100%  $\text{CH}_3\text{CN}$  over 25 minutes, 2.0 mL/min.
- NP-HPLC: retention time = 21.50 minutes (gradient from 100% Dichloromethane to 4:1 Dichloromethane: Methanol over 40 minutes)
- MS:  $\text{M}+\text{H} = 724$  by electrospray; calculated  $(\text{M}+\text{H}) = 724.8$ .

30

5

EXAMPLE 28N<sub>1</sub>, N<sub>2</sub>-(Cbz)<sub>2</sub>-S-methylisothiurea:

To a suspension of 10.02 g (36.0 mmol) of S-methylisothiurea in 100 mL of DCM was added 10 mL (40.0 mmol) of 4 M NaOH [prepared from 1.65 g NaOH and 10 mL of water]. The solid dissolved to produce a biphasic mixture and the flask was cooled in a wet ice/methanol bath. From one dropping funnel was added dropwise 20 mL (140 mmol) of benzyl chloroformate over a 15 minute period, simultaneously from a second dropping funnel was added 100 mL (100 mmol) of 1 M NaOH (prepared from 4.00 g NaOH and 100 mL of water) to maintain basicity of the biphasic mixture. After complete addition, the flask was allowed to warm to room temperature and the mixture stirred for 72 hrs.


The mixture was transferred to a separatory funnel and the layers separated. The aqueous phase was extracted with DCM. The combined organic extracts were washed twice with saturated brine, dried over MgSO<sub>4</sub>, and concentrated *in vacuo*. There was obtained 12.00 g (33.5 mmol, 93% yield) of product as a colorless oil which solidified upon storage at -20°C.

NP-HPLC: retention time = 11.38 minutes (isocratic elution using 1:1

Hexane: DCM)

NMR (CDCl<sub>3</sub>): 7.40 (s, C<sub>6</sub>H<sub>5</sub>); 5.33 (s, NH); 5.25 (s, OCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>); 2.50 (s, SCH<sub>3</sub>)

MS: M+H = 359 by DCI; calculated (M+H) = 359.4.

EXAMPLE 29Boc-D-Har(Cbz)<sub>2</sub>-OH:

A solution of 1.52 g (6.17 mmol) of Boc-D-Lys-OH, 2.33 g (6.53 mmol) of N<sub>1</sub>, N<sub>2</sub>-(Cbz)<sub>2</sub>-S-methylisothiurea, 5.0 mL (28.7 mmol) of DIEA, and 45 mL of methanol was stirred at room temperature under a N<sub>2</sub> atmosphere for 21 hrs.

The colorless solution was acidified to pH=1 with 10% HCl and extracted twice with ethyl acetate. The combined organic extracts were washed once with


SUBSTITUTE SHEET (RULE 26)

- 5 saturated brine, dried over  $\text{MgSO}_4$ , and concentrated *in vacuo*. Under high vacuum, there was obtained 2.82 g (5.07 mmol, 82% yield) of product as a colorless, viscous gel.

RP-HPLC: retention time = 13.76 minutes;  $\text{C}_{18}$ , O-100%  $\text{CH}_3\text{CN}$  over 25 minutes, 2.0 mL/min.

- 10 MS:  $\text{M}+\text{H} = 557$  by electrospray; calculated  $(\text{M}+\text{H}) = 557.6$ .

### EXAMPLE 30



#### Oxaloyl-D-Arg-Gly-Arg-H

- 15 Oxaloyl-D-Arg-Gly-Arg-H was prepared via solid phase synthesis according to the procedure of Webb *et al.* The Boc-Arg( $\text{NO}_2$ )-carbazone resin was prepared by loading  $\text{N}_\alpha$ -Boc( $\text{N}_\text{g}$ , $\text{NO}_2$ )argininal semicarbazonyl- $\text{N}_2$ -trans-4-methylenecyclohexane-1-carboxylic acid onto MBHA (methylbenzhydrylamine) resin (0.57 mmol/g) using the described procedure. Starting with 820 mg of Boc-
- 20 Arg( $\text{NO}_2$ )-carbazone resin, sequential peptide elongation was as follows: (1) the resin is treated with 8 mL of 40% TFA/ $\text{CH}_2\text{Cl}_2$  for 15 min. and washed 3 times with  $\text{CH}_2\text{Cl}_2$ , one time with 5% DIEA/ $\text{CH}_2\text{Cl}_2$ , and 3 times with  $\text{CH}_2\text{Cl}_2$ ; (2) coupling with 148 mg of Boc-Gly, 113 mg of HOBt, 318 mg HBTU, 292  $\mu\text{L}$  of DIEA in 10 mL DMF for 0.5 h; the resin is then washed once with DMF, 3 times with  $\text{CH}_2\text{Cl}_2$ , once
- 25 with MeOH, and 3 times with  $\text{CH}_2\text{Cl}_2$ ; (3) repeat step 1; (4) repeat step 2 using 229 mg of Boc-D-Arg( $\text{Cbz}$ ) $_2$ OH instead of Boc-Gly; (5) repeat step 1; (6) repeat step 2 using 150 mg of monobenzyloxalate; (7) the resin is then washed once with DMF, 3 times with  $\text{CH}_2\text{Cl}_2$ , once with MeOH, and 3 times with  $\text{CH}_2\text{Cl}_2$ . The protected peptide aldehyde is cleaved by shaking, the resin, a solution of 5 mL of THF, 1 mL
- 30 AcOH, 1 mL of 37% aqueous formaldehyde, and 0.1 mL of 1N HCl for 1h. The filtrate is collected and the resin is washed with 2 mL each of THF, 50% aqueous THF, and water. The combined filtrates are diluted with 25 mL of water and extracted 3 times with EtOAc. The combined organic layers are washed with 5%  $\text{NaHCO}_3$ , 2 times with water, once with brine, dried ( $\text{MgSO}_4$ ), and evaporated *in*
- 35 *vacuo* to give 78 mg of an oil: RP-HPLC retention: 12.04, 12.53, and 12.44 min


- 5 (tautomers). The oil in 2.5 mL of 10% aqueous MeOH containing 68  $\mu$ L AcOH and 50 mg of 5% palladium on charcoal (Degussa, 50% water by weight) is placed on a Parr hydrogenator and shaken for 2.25-3.5h at 30 psi. The solution is filtered and lyophilized to give 12.5 mg of a white solid.

RP-HPLC retention: 2.98 min; 3.47 min (thiosemicarbazone derivative); C18,

- 10 O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

MS: (M+H)= 444.5 by fast atom bombardment, calculated (M+H)= 444.2.

#### EXAMPLE 31



- 15 Malonoyl-D-Arg-Gly-Arg-H


Malonoyl-D-Arg-Gly-Arg-H was prepared via the solid phase method in Example 30 except that the growing peptide is capped with monobenzyloxymalonate. After lyophilization, a 40% yield of a white solid is obtained:

RP-HPLC retention: 2.98 min; 3.94 min (thiosemicarbazone derivative); C18,

- 20 O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

MS: (M+H)= 458.3 by fast atom bombardment, calculated (M+H)= 458.2.

#### EXAMPLE 32



- 25 Ph(CH<sub>2</sub>)<sub>2</sub>CO-(D-Arg)-Gly-Arg-H

3-Phenylpropanoyl-D-Arg-Gly-Arg-H was prepared in a similar fashion via the solid phase method in Example 30 except that the growing peptide is capped with 3-phenylpropanoic acid. The crude product after hydrogenolysis and filtration is purified by preparative RP-HPLC (18 mL/min, 2.5 X 25 cm Vydac C-18) using a linear gradient of 100% water/0.1% TFA to 100% CH<sub>3</sub>CN/0.1% TFA. The appropriate fractions are pooled and lyophilized to give a 57% yield of a white solid:


30

- 5 RP-HPLC retention: 8.65 min and 8.83 (tautomers); 8.87 min (thiosemicarbazone derivative); 11.02 min (2,4-dinitrophenylhydrazone derivative); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

MS: (M+H)= 504.3 by fast atom bombardment, calculated (M+H)= 504.3.

10

### EXAMPLE 33



#### BnSO<sub>2</sub>-D-Arg-Gly-Arg-H


- Benzylsulfonyl-D-Arg-Gly-Arg-H was prepared in a similar fashion via the solid phase method in Example 30 except that the growing peptide is capped with benzylsulfonyl chloride (6 eq) and DIEA (12 eq) in DCM. The crude, product after hydrogenolysis and filtration, is purified by preparative RP-HPLC (18 mL/min, 2.5 X 25 cm Vydac C-18) using a linear gradient of 100% water/0.1% TFA to 100% CH<sub>3</sub>CN/0.1% TFA. The appropriate fractions are pooled and lyophilized to give a 48% yield of a white solid:

- 20 RP-HPLC retention: 8.19 min and 8.51 (tautomers); 10.15 min (2,4-dinitrophenylhydrazone derivative); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

MS: (M+H)= 526.4 by fast atom bombardment, calculated (M+H)= 526.3.

25

### EXAMPLE 34



#### 2-NapOCH<sub>2</sub>CO-D-Arg-Gly-Arg-H

- 2-Napthoxyacetyl-D-Arg-Gly-Arg-H was prepared in a similar fashion via the solid phase method in Example 30 except that the arginine resin is protected as the bis-Cbz derivative made from the aldehyde of Example 27. The peptide was capped using (2-Naphthoxy)acetic acid N-hydroxysuccinimide ester. The crude product after hydrogenolysis and filtration is purified by preparative RP-HPLC (18


- 5 mL/min, 2.5 X 25 cm Vydac C-18) using a linear gradient of 100% water/0.1% TFA to 100% CH<sub>3</sub>CN/0.1% TFA. The appropriate fractions are pooled and lyophilized to give a 39% yield of a white solid:

RP-HPLC retention: 9.49 min and 9.74 (tautomers); 9.76 min (semicarbazone derivative); 11.57 min (2,4-dinitrophenylhydrazone derivative); C18,

- 10 O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

MS: (M+H)= 556.2 by fast atom bombardment, calculated (M+H)= 556.4.

#### EXAMPLE 35



- 15 Boc-D-Cit-Gly-Arg-H


Boc-D-Cit-Gly-Arg-H was prepared in a similar fashion via the solid phase method in Example 30. The peptide was capped with Boc-D-Citrulline. After cleavage and filtration from the resin, the filtrate is loaded directly onto a preparative RP-HPLC column (18 mL/min, 2.5 X 25 cm Vydac C-18) and eluted using a linear gradient of 100% water/0.1% TFA to 100% CH<sub>3</sub>CN/0.1% TFA. The appropriate fractions are pooled and lyophilized to give the protected peptide. The crude product after hydrogenolysis and filtration, is purified by preparative RP-HPLC (18 mL/min, 2.5 X 25 cm Vydac C-18) using a linear gradient of 100% water/0.1% TFA to 100% CH<sub>3</sub>CN/0.1% TFA. The appropriate fractions are pooled and lyophilized to give a 3.8% yield of a white solid:

RP-HPLC retention: 10.94 min and 10.56 (tautomers); 12.96 min

(thiosemicarbazone derivative); C18, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

MS: (M+H)= 473.3 by fast atom bombardment, calculated (M+H)= 473.4.

#### EXAMPLE 36




- 30

Boc-D-Har((CH<sub>3</sub>)<sub>4</sub>)-Gly-Arg-H

SUBSTITUTE SHEET (RULE 26)

- 5 A 2.3 mg (3.6  $\mu$ mol) sample of Boc-D-Lys-Gly-Arg-H, as prepared in Example 6, in 20  $\mu$ L of DMF is added 3.3 mg (8.6  $\mu$ mol) of HBTU followed by 1.6  $\mu$ L (10.8  $\mu$ mol) of DIEA. After 20h at 23°C, the reaction mixture is loaded directly onto a RP-HPLC column (2 mL/min, 0.1 X 25 cm Vydac C-18) and eluted using a linear gradient of 100% water/0.1% TFA to 100% CH<sub>3</sub>CN/0.1% TFA. The appropriate
- 10 fractions are pooled and lyophilized to give 1.6 mg (70%) of a white solid:
- RP-HPLC retention: 9.44 and 9.96 min (tautomers); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.
- MS: (M+H)= 542.4 by fast atom bombardment, calculated (M+H)= 542.4.


15

EXAMPLE 37Boc-D-Har-Gly-Arg-H

- Boc-D-Har-Gly-Arg-H was prepared in a similar fashion using the solid phase method of Example 30. The peptide was capped with Boc-Har-(Cbz<sub>2</sub>)-OH, as prepared in Example 29. The crude product after hydrogenolysis for 5h and
- 20 filtration is lyophilized to give a 32% yield of a white solid:
- RP-HPLC retention: 9.04 and 9.17 min (tautomers); 9.17 min (thiosemicarbazone derivative); 12.10 min (2,4-dinitrophenylhydrazone derivative); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.
- 25 MS: (M+H)= 486.3 by fast atom bombardment, calculated (M+H)= 486.4.




5

EXAMPLE 38Boc-D-Arg-Ala-Arg-H

Boc-D-Arg-Ala-Arg-H was prepared in a similar fashion via the solid phase method in Example 34 except that the growing peptide is coupled with Boc-Alanine instead of Boc-Glycine. The crude product after hydrogenolysis and filtration is purified by preparative RP-HPLC (18 mL/min, 2.5 X 25 cm Vydac C-18) using a linear gradient of 100% water/0.1% TFA to 100% CH<sub>3</sub>CN/0.1% TFA. The appropriate fractions are pooled and lyophilized to give a 15% yield of a white solid:

RP-HPLC retention: 11.24 min and 11.37 (tautomers); 16.88 min (2,4-dinitrophenylhydrazone derivative); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

MS: (M+H)= 486.3 by fast atom bombardment, calculated (M+H)= 486.3.

EXAMPLE 39

20

Boc-D-Arg-D-Ala-Arg-H

Boc-D-Arg-D-Ala-Arg-H was prepared in a similar fashion via the solid phase method in Example 34. The crude product after hydrogenolysis and filtration is lyophilized to give a white solid:

RP-HPLC retention: 11.53 min and 12.00 (tautomers); 11.71 min (thiosemicarbazone derivative); 16.83 min (2,4-dinitrophenylhydrazone derivative); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.


MS: (M+H)= 486.2 by fast atom bombardment, calculated (M+H)= 486.3.

30

EXAMPLE 40

SUBSTITUTE SHEET (RULE 26)

56



5

Boc-D-Arg-β-Ala-Arg-H


Boc-D-Arg-β-Ala-Arg-H was prepared in a similar fashion using the solid phase method in Example 34. The crude product after hydrogenolysis and filtration is purified by preparative RP-HPLC (18 mL/min, 2.5 X 25 cm Vydac C-18) using a linear gradient of 100% water/0.1% TFA to 100% CH<sub>3</sub>CN/0.1% TFA. The appropriate fractions are pooled and lyophilized to give a 52% yield of a white solid:

10

RP-HPLC retention: 11.48 min and 11.73 min (tautomers); 11.42 min (thiosemicarbazone derivative); 16.80 min (2,4-dinitrophenylhydrazone derivative); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

15

MS: (M+H)= 486.3 by fast atom bombardment, calculated (M+H)= 486.3.

EXAMPLE 41Boc-D-Arg-Aib-Arg-H

20

Boc-D-Arg-Aib-Arg-H was prepared in a similar fashion using the solid phase method in Example 34 except that the growing peptide is coupled with Boc-α-methylalanine instead of Boc-Gly and HATU is used as the coupling agent. The crude product after hydrogenolysis and filtration is purified by preparative RP-HPLC (18 mL/min, 2.5 X 25 cm Vydac C-18) using a linear gradient of 100% water/0.1% TFA to 100% CH<sub>3</sub>CN/0.1% TFA. The appropriate fractions are pooled and lyophilized to give a 23% yield of a white solid:

25

RP-HPLC retention: 11.84, 12.07 and 12.19 min (tautomers); 16.91 min (2,4-dinitrophenylhydrazone derivative); C<sub>18</sub>, O-100% CH<sub>3</sub>CN over 25 minutes, 2.0 mL/min.

30

5 MS: (M+H)= 500.2 by fast atom bombardment, calculated (M+H)= 500.3.

#### EXAMPLE 42

##### Boc-D-Arg(NO<sub>2</sub>)(OH)CONHEtPh

To a solution of Boc-Arg(NO<sub>2</sub>)(OH)COOH, prepared by the method of Webb,  
10 T. R. et al, Patent. No. WO 94/08941 (October 18, 1993), (850 mg, 2.4 mmol) and  
phenethylamine (609  $\mu$ L, 4.8 mmol) in 10 mL DMF is added BOP (1.3g, 2.9 mmol)  
followed by DIEA (630  $\mu$ L 3.6 mmol). The solution was allowed to stir for 18h. The  
reaction mixture was dissolved in EtOAc (250 mL) and washed with 1N HCl (75 mL),  
H<sub>2</sub>O (75 mL), 5% NaHCO<sub>3</sub> (75 mL) and brine (75 mL). The organic layer was dried  
15 over MgSO<sub>4</sub> and concentrated *in vacuo* to afford 900 mg (82%) of a light brown  
solid.

RP-HPLC: 15.30, 15.95 min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

MS:(M+H)= 453.2 by electrospray; calculated (M+H)= 453.8.

20

#### EXAMPLE 43

##### Boc-D-Arg(NO<sub>2</sub>)(OH)CONHBn

Boc-Arg(NO<sub>2</sub>)(OH)CONHBn was prepared according to Example 42 by  
coupling benzylamine to Boc-Arg(NO<sub>2</sub>)(OH)COOH in a 81% yield.

RP-HPLC: 14.6, 15.2 min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

25

MS:(M+H)= 439.0 by electrospray; calculated (M+H)= 439.4

#### EXAMPLE 44

##### Boc-D-Arg(NO<sub>2</sub>)(OH)CON(CH<sub>2</sub>)<sub>4</sub>

Boc-Arg(NO<sub>2</sub>)(OH)CON(CH<sub>2</sub>)<sub>4</sub> was prepared according to Example 42 by  
30 coupling pyrrolidine to Boc-Arg(NO<sub>2</sub>)(OH)COOH in a 44% yield.

RP-HPLC: 13.0, 13.1 min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

MS:(M+H)= 303.0 by electrospray (boc group removed in the spectrometer);  
calculated (M+H)= 403.4.

35

#### EXAMPLE 45

##### Boc-D-Arg(Cbz)<sub>2</sub>-Gly-OH

Boc-D-Arg(Cbz)<sub>2</sub>OH (Bachem) was coupled to glycine according to Example  
1 in quantitative yield as a white foam.

RP-HPLC: 18.9 min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

40

5

EXAMPLE 46BnSO<sub>2</sub>-D-Arg(Tos)-Gly-OH

To a solution of Boc-D-Arg(Tos)OH (11.2 g, 26.1 mmol), HOBt (3.9 g, 28.7 g), glycine benzyl ester hydrochloride (5.2 g, 26.1 mmol) in 52 mL of DCM is added DIEA (4.5 mL, 26.1 mmol) followed by DCC (5.9 g, 28.7 mmol; dissolved in 5 mL of DCM). After stirring for 18h, the reaction mixture is cooled to -10°C and the precipitate is filtered off. The filtrate is extracted twice with 1N HCl (50 mL), twice with 5% NaHCO<sub>3</sub> (50 mL), brine (50 mL), dried (MgSO<sub>4</sub>), and concentrated *in vacuo* to give 15.2 g (100%) of Boc-D-Arg(Tos)-Gly-OBn as a white solid. RP-HPLC: 18.9 min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

To a solution of D-Arg(Tos)-Gly-OBn (5.2 mmol; Boc previously removed by treatment with 40% TFA/DCM (0.1M) for 15 min. and then concentrated *in vacuo*; RP-HPLC: 13.8 min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min) in 21 mL of DCM is neutralized with DIEA before cooling to -78°C where 1.19 g (6.24 mmol) of BnSO<sub>2</sub>Cl is added as a solid followed by 900 µL (10.4 mmol). The solution is allowed to warm to 23°C and stir for 18h. The reaction mixture was extracted twice with 10% citric acid, twice with 5% NaHCO<sub>3</sub> (50 mL), brine (50 mL), dried (MgSO<sub>4</sub>), and concentrated *in vacuo* to give 3.2 g (97%) of BnSO<sub>2</sub>-D-Arg(Tos)-Gly-OBn as a white solid. RP-HPLC: 18.0 min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

The benzyl ester is removed by hydrogenation of 3.2 g of BnSO<sub>2</sub>-D-Arg(Tos)-Gly-OBn dissolved in 50 mL of MeOH with 800 mg of 5%Pd/C (50% wt. water), and 50 psi pressure of hydrogen on a Parr apparatus for 1.5h. The mixture is filtered through celite and concentrated *in vacuo* to give 2.54 g (93%) of the title compound as a white foam.

RP-HPLC: 15.0 min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

30

EXAMPLE 47Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)COOH

To a solution of 2.85 mg (4.75 mmol) of Boc-D-Arg(Cbz)<sub>2</sub>-Gly-OH and 0.66 g (5.7 mmol) of N-hydroxysuccinimide dissolved in 9.5 mL dioxane was added a solution of 1.08 g (5.23 mmol) of DCC dissolved in 4.0 mL of dioxane. The resultant solution was stirred at room temperature for 18h, during which time the DCU separated from the solution. The solids were removed by vacuum filtration and the clear colorless filtrate was treated with a single portion of Arg(NO<sub>2</sub>)(OH)COOH (1.658 g, 4.74 mmol, the Boc was previously removed with 40% TFA/DCM (0.1M) for 30 min, and concentrated *in vacuo*) and sodium bicarbonate (1.2 g, 14.22 mmol) dissolved in 8.5 mL of water. The resultant mixture was stirred at room temperature

- 5 18h. The solution was acidified to pH 1-2 with 10% HCl solution and extracted with ethyl acetate three times. The combined organics were washed twice with brine, dried over  $\text{MgSO}_4$ , and concentrated *in vacuo* to afford 2.4 g (54%) of a light brown solid.

RP-HPLC: 17.90 min;  $\text{C}_{18}$ , 0-100%  $\text{CH}_3\text{CN}$  over 25 min., 1.5 mL/min.

- 10 MS: (M+H)= 831.0; calculated (M+H)= 831.35.

#### EXAMPLE 48

##### Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CONHCH<sub>3</sub>

- To a solution of Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)COOH (200 mg, 0.22 mmol) and methylamine (9  $\mu\text{L}$ , 0.24 mmol) in 1 mL DMF, is added BOP (0.11 g, 0.26 mmol) followed by NMM (70  $\mu\text{L}$ , 0.65 mmol). The solution was allowed to stir 18h. The reaction mixture was dissolved in EtOAc (25 mL) and washed with 1N HCl (5 mL), Water (5 mL), 5%  $\text{NaHCO}_3$  (5 mL) and brine (5 mL). The organic layer was dried over  $\text{MgSO}_4$  and concentrated *in vacuo* to afford 145 mg (72%) as a light brown solid.

RP-HPLC: 17.80 min.;  $\text{C}_{18}$ , 0-100%  $\text{CH}_3\text{CN}$  over 25 min., 1.5 mL/min.

MS: (M+H)= 844.5; calculated (M+H)= 844.41.

#### EXAMPLE 49

##### 25 Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CONHPh

Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CONHPh was prepared according to Example 48 by coupling aniline to Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)COOH in a 80% yield.

RP-HPLC: 20.0 min.;  $\text{C}_{18}$ , 0-100%  $\text{CH}_3\text{CN}$  over 25 min., 1.5 mL/min.

- 30 MS: (M+H)= 906.6 by electrospray; calculated (M+H)= 906.5.

#### EXAMPLE 50

##### Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CONHEtPh

- To a solution of Boc-D-Arg(Cbz)<sub>2</sub>-Gly-OH (771 mg, 1.29 mmol) in 5 mL of DMF was added a solution of Arg(NO<sub>2</sub>)(OH)CONHEtPh, (Boc was previously removed with 40% TFA/DCM (0.1M) for 30 min, and dried *in vacuo* in DMF (2.15 mL, reaction conc 0.2M) and BOP (695 mg, 1.57 mmol) followed by DIEA (746  $\mu\text{L}$ , 4.29 mmol). The solution was allowed to stir 18h. The reaction mixture was dissolved in EtOAc (250 mL) and washed with 1N HCl (75 mL),  $\text{H}_2\text{O}$  (75 mL), 5%  $\text{NaHCO}_3$  (75 mL) and brine (75 mL). The organic layer was dried over  $\text{MgSO}_4$  and concentrated *in vacuo* to afford 1.19 g (89%) as an off white solid.

- 5            RP-HPLC: 19.40 min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.  
             MS:(M+H)= 934.3; calculated (M+H)= 934.3

#### EXAMPLE 51

##### BnSO<sub>2</sub>-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CONHEtPh

- 10            To a 0°C solution of D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CONHEtPh (990 mg, 1.1 mmol; Boc was previously removed with 40%TFA/ DCM (0.1M) for 30 min, and concentrated *in vacuo*) in DCM (4.24 mL) and DIEA (555  $\mu$ L, 3.3 mmol) was added BnSO<sub>2</sub>Cl (222 mg, 1.2 mmol). The solution was allowed to stir 18h after which the reaction mixture was dissolved in EtOAc (250 mL) and washed with 1N HCl (75 mL),  
15            H<sub>2</sub>O (75 mL), 5% NaHCO<sub>3</sub> (75 mL) and brine (75 mL). The organic layer was dried over MgSO<sub>4</sub> and concentrated *in vacuo* to afford 440 g (42%) as an off-white solid.  
             RP-HPLC: 19.79 min. C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.  
             MS:(M+H)= 988.4; calculated (M+H)= 988.1.

- 20            EXAMPLE 52

##### BnSO<sub>2</sub>-D-Arg-Gly-Arg-CONHEtPh

- To BnSO<sub>2</sub>-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CONHEtPh (200 mg, 0.202 mmol) was added Anisole (1 mL), ethylmethylsulfide (0.5 mL), and HF (20 mL anhydrous). The reaction was allowed to stir for 1.5h at 0°C. The HF was removed *in vacuo* and  
25            the reaction mixture washed 3 times with chilled ether, glacial acetic acid, and water. The aqueous wash was lyophilized directly and 100 mg (0.148 mmol) of this crude material redissolved in 1M TFA in DMSO (222  $\mu$ L, 1.5eq) and IBX (1.48 mL, 0.5M in DMSO). The reaction is allowed to stir 18h. The reaction mixture is purified directly on a 5.0 cm C<sub>18</sub> column and lyophilized directly to give 74.1 mg (74%) as a  
30            fluffy white solid.  
             RP-HPLC: 12.96 min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.  
             MS:(M+H)= 673.2; calculated (M+H)= 673.3.

#### EXAMPLE 53

##### Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)CONHEtPh

- To a solution of Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CONHEtPh (275 mg, 0.294 mmol) in DMSO/Toluene(1:1) (6 mL) was added EDC (731 mg, 3.8 mmol) and DCA (97  $\mu$ L, 1.18 mmol). The reaction was allowed to stir for 18h. The reaction was purified directly by flash silica gel chromatography(10% MeOH/DCM/0.1% AcOH) to  
40            yield 190 mg (69%) of a white solid.

5            RP-HPLC: 19.27 min.(2 peaks); C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

MS:(M+H)= 932.5; calculated (M+H)= 932.28

#### EXAMPLE 54

##### 10    Boc-D-Arg-Gly-Arg-CONHEtPh

To a solution of Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)CONHEtPh (190 mg, 0.204 mmol) in MeOH (15.4 mL) was added 10% Pd/C (153 mg, 70mol%) and 1N HCl (402 uL) and H<sub>2</sub> at 50 psi. The reaction was allowed shake for 8h. The catalyst was removed by filtration through celite and the reaction mixture diluted with water and lyophilized. The crude solid was redissolved in water and purified directly on a 2.5  
15    cm x 25 cm C<sub>18</sub> column (0-60% ACN/0.1% TFA) to yield 42 mg (33%) of a white solid.

RP-HPLC: 13.44 min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

MS:(M+H)= 619.3; calculated (M+H)= 619.3

20

#### EXAMPLE 55

##### D-Arg-Gly-Arg-CONHEtPh

To Boc-D-Arg-Gly-ArgCONHEtPh (2.3 mg, 0.004 mmol) was added 40%TFA/DCM (0.1M) for 30 min, and dried *in vacuo*.

25            RP-HPLC: 11.77 min. C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

MS:(M+H)= 519.0; calculated (M+H)= 519.3

#### EXAMPLE 56

##### Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CONHBn

30            Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CONHBn was prepared according to Example 47 by coupling Arg-(NO<sub>2</sub>)(OH)CONHBn to Boc-D-Arg(Cbz)<sub>2</sub>-Gly-OH in a 60% yield.

RP-HPLC: 19.32 min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

MS:(M+H)= 920.5 by electrospray; calculated (M+H)= 921.0.

35

#### EXAMPLE 57

##### Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CON(CH<sub>2</sub>)<sub>4</sub>

Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CON(CH<sub>2</sub>)<sub>4</sub> was prepared according to Example 50 by coupling Arg-(NO<sub>2</sub>)(OH)CON(CH<sub>2</sub>)<sub>4</sub> to Boc-D-Arg(Cbz)<sub>2</sub>-Gly-OH. The  
40    crude organic layer, after workup, was purified by flash silica gel chromatography (8% MeOH/DCM/0.1% AcOH) to yield 117 mg (59%) of a white solid.

5 RP-HPLC: 17.96 min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.  
MS:(M+H)= 884.5 electrospray; calculated (M+H)= 883.98.

### EXAMPLE 58

BnSO<sub>2</sub>-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CON(CH<sub>2</sub>)<sub>4</sub>

10  $\text{BnSO}_2\text{-D-Arg(Cbz)}_2\text{-Gly-Arg(NO}_2\text{)(OH)CON(CH}_2\text{)}_4$  was prepared according to Example 51. The reaction mixture was purified directly by flash silica gel chromatography (10% MeOH/DCM/0.1% AcOH) to yield 35 mg (37%).

RP-HPLC: 18.12min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

### EXAMPLE 59

BnSO<sub>2</sub>-D-Arg-Gly-Arg-CON(CH<sub>2</sub>)<sub>4</sub>

BnSO<sub>2</sub>-D-Arg-Gly-ArgCON(CH<sub>2</sub>)<sub>4</sub> was prepared according to Example 52 to yield 2.3 mg (14%).

RP-HPLC: 12.09 C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

20 MS:(M+H)= 623.2 electrospray; calculated (M+H)= 623.3.

### EXAMPLE 60

**Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)CONHBn**

Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CONHBn was prepared according to  
25 Example 53. The reaction mixture was purified directly by flash silica gel  
chromatography(10%MeOH/DCM/.1% AcOH) to yield 48.5 mg (32%).

RP-HPLC: 19.80 (2 peaks); C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

MS:(M+H)= 918.4 by electrospray; calculated (M+H)= 918.81.

### EXAMPLE 61

**Boc-D-Arg-Gly-Arg-CONHBn**

Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CONHBn was oxidized and purified as in Example 53 followed by hydrogenation and purification as in Example 13 to yield 5 mg (17%).

35 RP-HPLC: 13.76 broad peak; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

MS:(M+H)= 605.5 electrospray; calculated (M+H)= 605.7.

### EXAMPLE 62

Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)CONHCH<sub>3</sub>



- 5        Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CONHCH<sub>3</sub> was prepared according to Example 53. The reaction mixture was diluted with EtOAc and washed with water, brine, and dried over MgSO<sub>4</sub> to yield 47 mg (39%).

RP-HPLC: 17.85(2 peaks) C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

10

EXAMPLE 63Boc-D-Arg-Gly-Arg-CONHCH<sub>3</sub>

Boc-Arg-Gly-Arg-CONHCH<sub>3</sub> was prepared according to Example 54 to yield 5 mg (17%).

RP-HPLC: 13.76 broad peak C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

15

MS:(M+H)= 529.0 electrospray; calculated (M+H)= 529.32

EXAMPLE 64Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)CONHPh

- 20        Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)(OH)CONHPh was oxidized according to Example 53. The reaction mixture was diluted with EtOAc and washed with water, brine, and dried over MgSO<sub>4</sub> to yield 96 mg (96%) of a white foam.

RP-HPLC: 20.0; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

MS:(M+H)= 904.5.0 by electrospray; calculated (M+H)= 904.3.

25

EXAMPLE 65D-Arg-Gly-Arg-CONHPh

Boc-D-Arg(Cbz)<sub>2</sub>-Gly-Arg(NO<sub>2</sub>)CONHPh was submitted to the HF cleavage conditions of Example 11 to yield 2 mg (18.4%) of a white solid.

RP-HPLC: 11.42 broad peak C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

30

MS:(M+H)= 491.0 electrospray; calculated (M+H)= 491.6

EXAMPLE 66Boc-D-Arg(Tos)-Gly-Arg(NO<sub>2</sub>)(OH)CONBn

- 35        The titled compound was prepared according to Example 50 by coupling Arg-(NO<sub>2</sub>)(OH)CONHBn to BnSO<sub>2</sub>-D-Arg(Tos)-Gly-OH except that upon dilution of the organic layer and addition of 1N HCl, a oily precipitate is formed which is collected and re-precipitated with MeOH and water. The solid is collected, redissolved in MeOH and lyophilized to give a 38% yield of a white solid.

RP-HPLC: 15.9 min.; C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.

40

EXAMPLE 67

5 BnSO<sub>2</sub>-D-Arg-Gly-Arg-CONBn

BnSO<sub>2</sub>-D-Arg-Gly-ArgCONBn was prepared according to Example 52 to yield 110 mg (73%) of a fluffy white solid.

RP-HPLC: 11.6 min (broad peak) C<sub>18</sub>, 0-100% CH<sub>3</sub>CN over 25 min., 1.5 mL/min.  
MS:(M+H)= 659.2 electrospray; calculated (M+H)= 659.3.

10

EXAMPLE 68

Evaluation of the compounds of this invention is guided by *in vitro* protease activity assays (see below) and *in vivo* studies to evaluate biological half-life, antithrombotic efficacy, and effects on hemostasis and hematological parameters (see Example 69 below).

The compounds of the present invention are dissolved in buffer to give solutions containing concentrations such that assay concentrations range from 0 to 100 µM. In the assays for thrombin, prothrombinase and factor Xa, a synthetic chromogenic substrate is added to a solution containing test compound and the enzyme of interest and the residual catalytic activity of that enzyme is determined spectrophotometrically. The IC<sub>50</sub> of a compound is determined from the substrate turnover. The IC<sub>50</sub> is the concentration of test compound giving 50% inhibition of the substrate turnover. The compounds of the present invention desirably have an IC<sub>50</sub> of less than 500 nM in the factor Xa assay, preferably less than 200 nM, and more preferred compounds have an IC<sub>50</sub> of about 100 nM or less in the factor Xa assay. The compounds of the present invention desirably have an IC<sub>50</sub> of less than 4.0 µM in the prothrombinase assay, preferably less than 200 nM, and more preferred compounds have an IC<sub>50</sub> of about 10 nM or less in the prothrombinase assay. The compounds of the present invention desirably have an IC<sub>50</sub> of greater than 1.0 µM in the thrombin assay, preferably greater than 10.0 µM, and more preferred compounds have an IC<sub>50</sub> of greater than 100.0 µM in the thrombin assay.

Amidolytic Assays for determining protease inhibition activity

The factor Xa and thrombin assays were performed at room temperature, in 0.02 M Tris-HCl buffer, pH 7.5, containing 0.15 M NaCl. The rates of hydrolysis of the para-nitroanilide substrate S-2765 (Chromogenix) for factor Xa, and the substrate Chromozym TH (Boehringer Mannheim) for thrombin following preincubation of the enzyme with inhibitor for 5 minutes at room temperature, and were determined using the Softmax 96-well plate reader (Molecular Devices), monitored at 405 nm to measure the time dependent appearance of p-nitroaniline.

5           The prothrombinase inhibition assay was performed in a plasma free system with modifications to the method described by Sinha, U. *et al.*, *Thromb. Res.*, 75, 427-436 (1994). Specifically, the activity of the prothrombinase complex was determined by measuring the time course of thrombin generation using the p-nitroanilide substrate Chromozym TH. The assay consists of preincubation ( 5  
10 minutes) of selected compounds to be tested as inhibitors with the complex formed from factor Xa (0.5 nM), factor Va (2 nM), phosphatidyl serine:phosphatidyl choline (25:75, 20 µM) in 20 mM Tris-HCl buffer, pH 7.5, containing 0.15 M NaCl, 5 mM CaCl<sub>2</sub> and 0.1% bovine serum albumin. Aliquots from the complex-inhibitor mixture were added to prothrombin (1 nM) and Chromozym TH (0.1 mM). The rate of  
15 substrate cleavage was monitored at 405 nm for two minutes. Eight different concentrations of inhibitor were assayed in duplicate. A standard curve of thrombin generation by an equivalent amount of untreated complex was used for determination of percent inhibition.

20

#### EXAMPLE 69

A series of studies were accomplished in rabbits to evaluate the biological half-life, antithrombotic efficacy, and effects on hemostasis and hematological parameters of the compound Boc-D-Arg-Gly-Arg-H (referred to in this example as "Boc-RGR-H").

#### Pharmacokinetics/Pharmacodynamics

25           Rabbits were anesthetized with intramuscular (I.M.) injections of Ketamine, Xylazine, and Acepromazine cocktail. A marginal ear vein and femoral vein were cannulated for drug administration and blood sampling. Citrated blood samples were obtained serially for two hours after intravenous (I.V.) bolus injection of Boc-RGR-H (2.0 mg/kg). Coagulation parameters [Activated Partial Thromboplastin times (aPTT),  
30 Prothrombin times (PT), and Fibrinogen (FIB)] and hematological parameters [ red blood cell (RBC), white blood cell (WBC), hematocrit (HcT), and platelet (PLT) counts] were measured on samples obtained at designated time points. Plasma concentrations of Boc-RGR-H were determined by HPLC.

#### Pk/Pd Results

35           Boc-RGR-H (2.0 mg/kg) exhibited a plasma concentration-dependent anticoagulative effect. The peak plasma concentration by HPLC was 3.78 µg/mL. This dose of Boc-RGR-H generated a 2.45 fold aPTT prolongation as compared to pretreatment values. The PT value was also prolonged by 1.15 fold and FIB were slightly prolonged 1.07 fold. There were no significant changes in hematological parameters as  
40 compared to saline controls.

5           Following I.V. administration, Boc-RGR-H was cleared from the plasma in a  
monoexponential decay (one-compartment model). The mean plasma half-life of Boc-  
RGR-H in rabbits was 70.6 minutes (n=3). The mean clearance value (CL) was 1.16  
mL/min/kg with a volume of distribution of 118.3 mL/kg. Dosing rate for infusion studies  
was determined by {Dosing rate = CL x C<sub>ss</sub>} where CL is clearance and C<sub>ss</sub> is plasma  
10           concentration at steady state.

#### Antithrombotic Efficacy in a Rabbit Model of Venous Thrombosis

          A rabbit deep vein thrombosis model as described by Hollenbach, S. et al.,  
Thromb. Haemost. 71, 357-362 (1994), was used to determine the in-vivo antithrombotic  
activity of the test compounds. Rabbits were anesthetized with I.M. injections of  
15           Ketamine, Xylazine, and Acepromazine cocktail. A standardized protocol consisted of  
insertion of a thrombogenic cotton thread and copper wire apparatus into the abdominal  
vena cava of the anesthetized rabbit. A non-occlusive thrombus was allowed to develop  
in the central venous circulation and inhibition of thrombus growth was used as a  
measure of the antithrombotic activity of the studied compounds. Test agents or control  
20           saline were administered through a marginal ear vein catheter. A femoral vein catheter  
was used for blood sampling prior to and during steady state infusion of test compound.  
Initiation of thrombus formation begins immediately after advancement of the cotton  
thread apparatus into the central venous circulation. Test compounds were administered  
from time = 30 min to time = 150 min at which the experiment was terminated. The  
25           rabbits were euthanized and the thrombus excised by surgical dissection and  
characterized by weight and histology. Blood samples were analyzed for plasma  
concentrations and changes in hematological and coagulation parameters.

#### Effects of Boc-RGR-H in Rabbit Venous Thrombosis model

          Administration of Boc-RGR-H in the rabbit venous thrombosis model  
30           demonstrated antithrombotic efficacy at the higher doses evaluated. There was a dose-  
dependent effect of the compound on the aPTT and PT prolongation with the highest  
dose (1500 µg/kg + 14.85 µg/kg/min) extending aPTT and PT, 1.61 and 1.08 fold  
respectively (see Table 2). Boc-RGR-H had no significant effects on hematological  
35           parameters as compared to saline controls (see Table 3).

**TABLE 2 - ANTITHROMBOTIC EFFECTS OF Boc-RGR-H IN RABBITS**

	Dose Regimen		% Inhibition of Thrombosis	--fold increase over baseline--	
	(µg/kg + µg/kg/min)	n#		aPTT	PT
40	saline control	7	0.0	1.00 ± 0.05	1.01 ± 0.00
	200 + 1.98	5	8.4	1.04 ± 0.05	1.02 ± 0.01

5	600 + 5.94	5	-6.6	$1.15 \pm 0.02$	$1.03 \pm 0.01$
	1000 + 9.90	5	58.6	$1.31 \pm 0.04$	$1.06 \pm 0.01$
	1500 + 14.85	5	31.4	$1.61 \pm 0.13$	$1.08 \pm 0.01$

All measurements are an average of all samples after steady state administration of vehicle or Boc-RGR-H. Values are expressed as mean  $\pm$  SD.

10

TABLE 3 - EFFECTS OF Boc-RGR-H ON HEMATOLOGICAL PARAMETERS

Dose Regimen ( $\mu\text{g/kg}$ + $\mu\text{g/kg/min}$ )	n#	RBC $\times 10^6/\mu\text{L}$	WBC $\times 10^3/\mu\text{L}$	PLT $\times 10^3/\mu\text{L}$	Hct %
saline control	7	$5.64 \pm 0.49$	$4.30 \pm 1.63$	$432 \pm 129$	$35.2 \pm 2.81$
15 200 + 1.98	5	$5.69 \pm 0.14$	$4.45 \pm 0.97$	$393 \pm 84$	$36.9 \pm 3.90$
600 + 5.94	5	$5.74 \pm 0.48$	$3.87 \pm 0.65$	$442 \pm 81$	$35.3 \pm 3.01$
1000 + 9.90	5	$5.71 \pm 0.28$	$5.09 \pm 1.61$	$375 \pm 73$	$35.5 \pm 1.01$
1500 + 14.85	5	$5.80 \pm 0.50$	$4.82 \pm 1.01$	$433 \pm 22$	$36.0 \pm 3.31$

All measurements are an average of samples after steady state administration of vehicle

20 or

Boc-RGR-H. Values are mean  $\pm$  SD.

#### Effects of Boc-RGR-H on Cuticle Bleeding times in Rabbits

In an ancillary study, a Cuticle Bleeding Time model was used to evaluate the effect of test compounds on bleeding time (BT) measurements in rabbits. The standardized protocol involved anesthesia and catheterization of rabbits as previously described in this Example 43. To measure BT, a toe nail was cut 5mm from the base of the cuticle with a canine nail trimming device. Time until cessation of bleeding was recorded by wicking the blood on to BT blotting paper in 30 second intervals. Test compound was administered as a bolus followed by infusion regimen. Two BT measurements were obtained prior to administration of test compound and three BT measurements accomplished serially after initiation of steady state infusion of test compound. BT measurements were allowed to continue for up to 30 minutes at which time it was stopped with application of silver nitrate. Blood samples were obtained just prior to all BT measurements and processed for measurement of hematological and coagulation parameters.


The effect of Boc-RGR-H on rabbit cuticle BT at a dose of 1500  $\mu\text{g/kg}$  + 14.85  $\mu\text{g/kg/min}$  was a increase of 2.14 fold ( $\pm 1.26$ , n=4) from pre-administration baseline measurements. In these experiments average increase in aPTT and PT was 1.61 and 1.07 fold respectively. There were no significant changes in hematological parameters from pre to post Boc-RGR-H administration.

5           Although the invention has been described with reference to the disclosed  
embodiments, those skilled in the art will readily appreciate that the specific  
experiments detailed are only illustrative of the invention. It should be understood  
that various modifications can be made without departing from the spirit of the  
invention. Accordingly, the invention is limited only by the following claims.

10

5 WHAT IS CLAIMED IS:

1. A compound represented by the formula:



10 wherein:

$$m = 0, 1, 2, 3, 4;$$
$$n = 0, 1, 2, 3, 4;$$

$p = 0, 1, 2, 3, 4;$

$$q = 0, 1, 2, 3, 4;$$

15 Y = CHO, COCF<sub>3</sub>, COCF<sub>2</sub>CF<sub>3</sub>, COCO<sub>2</sub>R<sub>7</sub>, COCONR<sub>8</sub>R<sub>9</sub>, B(OR<sub>10</sub>)<sub>2</sub>;

where: R7, R8, R9, R10 are the same or different and = H, C1-

6alkyl, C3-8cycloalkyl, arylC<sub>1-3</sub>alkyl, heteroarylC<sub>1-3</sub>alkyl;

A = piperdinyl, pyrrolidinyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, C<sub>3-6</sub>heteroaryl, or is absent;

20  $R_1 = \text{H or C}_{1-3}\text{alkyl:}$

$J = 0$  or  $H_2$ :

**R<sub>2</sub> = H or C<sub>1-3</sub>alkyl;**

$$D = N, CH, NCH_2, NCH_2CH_2, CHCH_2;$$

**R<sub>3</sub> = H or C<sub>1-3</sub>alkyl;**

25 E = O or H<sub>2</sub>;

$R_4 = \text{H or CH}_3$ ;


**Q =** piperdinyl, pyrrolidinyl, C3-8 cycloalkyl, phenyl, substituted phenyl, naphthyl, pyridyl, or is absent;

**G = N, CH, or is H;**

30 R<sub>5</sub> = H or C<sub>1-3</sub> alkyl, or is absent if G is H;

- 5  $R_6 =$  H or  $CH_3$ ;
- $W =$  H, arylacyl, heteroarylacyl, arylC<sub>1-3</sub>alkylsulfonyl, arylsulfonyl, substituted arylsulfonyl, arylC<sub>1-4</sub>alkenylsulfonyl, C<sub>1-8</sub> alkylsulfonyl, heteroarylC<sub>1-3</sub>alkylsulfonyl, heteroarylsulfonyl, aryloxycarbonyl, C<sub>1-6</sub> alkyloxycarbonyl, arylC<sub>1-3</sub>alkyloxycarbonyl, arylaminocarbonyl, C<sub>1-6</sub> alkylaminocarbonyl, arylC<sub>1-3</sub>alkylaminocarbonyl,  $HOOC-C_0$ -3alkylcarbonyl, or is absent if G is H;
- 10  $X =$  H, C<sub>1-3</sub>alkyl,  $NR'R''$ ,  $NH-C(NR'R'')=NH$ ,  $NH-C(NHR')=NR''$ ,  $S-C(NR'R'')=NH$ ,  $S-C(NHR')=NR''$ ,  $C(NR'R'')=NH$ ,  $C(NHR')=NR''$ , or  $CR'=NR''$ ; where:  $R', R''$  are the same or different and  $=$  H, C<sub>1-6</sub>alkyl, C<sub>1-3</sub>arylalkyl, aryl or where  $R'R''$  forms a cyclic ring containing  $(CH_2)_p$  where  $p=2-5$ , with the proviso that when X is H or C<sub>1-3</sub>alkyl, then A must contain at least one N atom;
- 15  $Z =$  H, C<sub>1-3</sub>alkyl,  $NR'R''$ ,  $NH-C(NR'R'')=NH$ ,  $NH-C(NHR')=NR''$ ,  $S-C(NR'R'')=NH$ ,  $S-C(NHR')=NR''$ ,  $C(NR'R'')=NH$ ,  $C(NHR')=NR''$ , or  $CR'=NR''$ ; where:  $R', R''$  are the same or different and  $=$  H, C<sub>1-6</sub>alkyl, C<sub>1-3</sub>arylalkyl, aryl or where  $R'R''$  forms a cyclic ring containing  $(CH_2)_p$  where  $p=2-5$ , with the proviso that when Z is H or C<sub>1-3</sub>alkyl, then Q must contain at least one N atom;
- 20 and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.
- 25

2. The compound of claim 1, having the formula:





5 wherein:

m = 0,1,2,3,4;

n = 0,1,2,3,4;

p = 0,1,2,3,4;

q = 0,1,2,3,4;

10 Y = CHO, COCF<sub>3</sub>, COCF<sub>2</sub>CF<sub>3</sub>, COCO<sub>2</sub>R<sub>7</sub>, COCONR<sub>8</sub>R<sub>9</sub>; where:

R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub> are the same or different and = H, C<sub>1-6</sub>alkyl, C<sub>3-8</sub>cycloalkyl, arylC<sub>1-3</sub>alkyl, heteroarylC<sub>1-3</sub>alkyl;

A = piperdiny, pyrrolidiny, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, C<sub>3-6</sub>heteroaryl or is absent;

15 R<sub>2</sub> = H or C<sub>1-3</sub>alkyl;

Q = piperdiny, pyrrolidiny, C<sub>3-8</sub> cycloalkyl, phenyl, substituted phenyl, naphthyl, pyridyl, or is absent;

W = H, arylacyl, heteroarylacyl, arylC<sub>1-3</sub>alkylsulfonyl, arylsulfonyl, substituted arylsulfonyl, arylC<sub>1-4</sub>alkenylsulfonyl, C<sub>1-8</sub> alkylsulfonyl, heteroarylC<sub>1-3</sub>alkylsulfonyl, heteroarylsulfonyl, aryloxycarbonyl, C<sub>1-6</sub> alkyloxycarbonyl, arylC<sub>1-3</sub>alkyloxycarbonyl, arylaminocarbonyl, C<sub>1-6</sub>alkylaminocarbonyl, arylC<sub>1-3</sub>alkylaminocarbonyl, or HOOC-C<sub>0-3</sub>alkylcarbonyl;

20 X = H, C<sub>1-3</sub>alkyl, NR'R'', NH-C(NR'R'')=NH, NH-C(NHR')=NR'', S-C(NR'R'')=NH, S-C(NHR')=NR'', C(NR'R'')=NH, C(NHR')=NR'', or CR'=NR''; where: R', R'' are the same or different and = H, C<sub>1-6</sub>alkyl, C<sub>1-3</sub>arylalkyl, aryl or where R'R'' forms a cyclic ring containing (CH<sub>2</sub>)<sub>p</sub> where p=2-5, with the proviso that when X is H or C<sub>1-3</sub>alkyl, then A must contain at least one N atom;

25 Z = H, C<sub>1-3</sub>alkyl, NR'R'', NH-C(NR'R'')=NH, NH-C(NHR')=NR'', S-C(NR'R'')=NH, S-C(NHR')=NR'', C(NR'R'')=NH, C(NHR')=NR'', or CR'=NR''; where: R', R'' are the same or different and = H, C<sub>1-6</sub>alkyl, C<sub>1-3</sub>arylalkyl, aryl or where R'R'' forms a cyclic ring containing (CH<sub>2</sub>)<sub>p</sub> where p=2-5, with the proviso that when Z is H or C<sub>1-3</sub>alkyl, then Q must contain at least one N atom;

30 and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

- 5     3.     The compound of claim 1, having an IC<sub>50</sub> for Factor Xa of less than about 200 nM.
4.     The compound of claim 1, having an IC<sub>50</sub> for prothrombinase of less than about 2.0 μM.
- 10     5.     The compound of claim 1, having an IC<sub>50</sub> for thrombin of greater than about 1.0 μm.
6.     A compound selected from a group consisting of:
- 15     H-D-Arg-Gly-Arg-H  
Boc-D-Arg-Gly-Arg-H  
HOCCO-D-Arg-Gly-Arg-H  
HOCCCH<sub>2</sub>CO-D-Arg-Gly-Arg-H  
HOOC(CH<sub>2</sub>)<sub>2</sub>CO-D-Arg-Gly-Arg-H
- 20     PhCH<sub>2</sub>CH<sub>2</sub>CO-D-Arg-Gly-Arg-H  
PhCH<sub>2</sub>SO<sub>2</sub>-D-Arg-Gly-Arg-H  
EtOCO-D-Arg-Gly-Arg-H  
2-NaphthoxyAc-D-Arg-Gly-Arg-H  
Boc-D-Cit-Gly-Arg-H
- 25     Boc-D-Lys-Gly-Arg-H  
Boc-D-Har-Gly-Arg-H  
Boc-D-Har((CH<sub>3</sub>)<sub>4</sub>)-Gly-Arg-H  
Boc-D-Arg-Ala-Arg-H  
Boc-D-Arg-D-Ala-Arg-H
- 30     Boc-D-Arg-β-Ala-Arg-H  
Boc-D-Arg-Aib-Arg-H  
Boc-D-(2,3-Dap)-Gly-Arg-H  
Boc-D-(2,4-Dab)-Gly-Arg-H  
γ-Abu-Gly-Arg-H
- 35     Boc-D-Orn-Gly-Arg-H  
Boc-D-homoLys-Gly-Arg-H  
Boc-Bag-Gly-Arg-H  
Boc-D-4-Gpa-Gly-Arg-H

- 5        Boc-D-3-Gpa-Gly-Arg-H  
          Boc-D-4-Apa-Gly-Arg-H  
          Boc-D-3-Apa-Gly-Arg-H  
          Boc-D-4-Acg-Gly-Arg-H  
          Boc-D-(4-NH<sub>2</sub>Phe)-Gly-Arg-H
- 10        Boc-D-(3-NH<sub>2</sub>Phe)-Gly-Arg-H  
          BnSO<sub>2</sub>-D-(2,3-Dap)-Gly-Arg-H  
          BnSO<sub>2</sub>-D-(2,4-Dab)-Gly-Arg-H  
          BnSO<sub>2</sub>-D-Om-Gly-Arg-H  
          BnSO<sub>2</sub>-D-homoLys-Gly-Arg-H
- 15        BnSO<sub>2</sub>-Bag-Gly-Arg-H  
          BnSO<sub>2</sub>-D-4-Gpa-Gly-Arg-H  
          BnSO<sub>2</sub>-D-3-Gpa-Gly-Arg-H  
          BnSO<sub>2</sub>-D-4-Apa-Gly-Arg-H  
          BnSO<sub>2</sub>-D-3-Apa-Gly-Arg-H
- 20        BnSO<sub>2</sub>-D-4-Acg-Gly-Arg-H  
          BnSO<sub>2</sub>-D-(4-NH<sub>2</sub>Phe)-Gly-Arg-H  
          BnSO<sub>2</sub>-D-(3-NH<sub>2</sub>Phe)-Gly-Arg-H  
          BnSO<sub>2</sub>-D-(2,3-Dap)-Gly-Arg-CONH<sub>2</sub>  
          BnSO<sub>2</sub>-D-(2,4-Dab)-Gly-Arg-CONH<sub>2</sub>
- 25        BnSO<sub>2</sub>-D-Om-Gly-Arg-CONH<sub>2</sub>  
          BnSO<sub>2</sub>-D-homoLys-Gly-Arg-CONH<sub>2</sub>  
          BnSO<sub>2</sub>-Bag-Gly-Arg-CONH<sub>2</sub>  
          BnSO<sub>2</sub>-D-4-Gpa-Gly-Arg-CONH<sub>2</sub>  
          BnSO<sub>2</sub>-D-3-Gpa-Gly-Arg-CONH<sub>2</sub>
- 30        BnSO<sub>2</sub>-D-4-Apa-Gly-Arg-CONH<sub>2</sub>  
          BnSO<sub>2</sub>-D-3-Apa-Gly-Arg-CONH<sub>2</sub>  
          BnSO<sub>2</sub>-D-4-Acg-Gly-Arg-CONH<sub>2</sub>  
          BnSO<sub>2</sub>-D-(4-NH<sub>2</sub>Phe)-Gly-Arg-CONH<sub>2</sub>  
          BnSO<sub>2</sub>-D-(3-NH<sub>2</sub>Phe)-Gly-Arg-CONH<sub>2</sub>
- 35        BnSO<sub>2</sub>-D-Arg-Gly-(2,4-Dab)-H  
          BnSO<sub>2</sub>-D-Arg-Gly-(homoLys)-H  
          BnSO<sub>2</sub>-D-Arg-Gly-(4-Gpa)-H

- 5            BnSO<sub>2</sub>-D-Arg-Gly-(3-Gpa)-H  
             BnSO<sub>2</sub>-D-Arg-Gly-(4-Apa)-H  
             BnSO<sub>2</sub>-D-Arg-Gly-(3-Apa)-H  
             BnSO<sub>2</sub>-D-Arg-Gly-(4-Acg)-H  
             BnSO<sub>2</sub>-D-Arg-Gly-(4-NH<sub>2</sub>Phe)-H
- 10           BnSO<sub>2</sub>-D-Arg-Gly-(3-NH<sub>2</sub>Phe)-H  
             Boc-D-Arg-Gly-Arg-CONHEtPh  
             Boc-D-3-PIA-Gly-Arg-CONHEtPh  
             Boc-D-4-PIA-Gly-Arg-CONHEtPh  
             Boc-D-3-GPIA-Gly-Arg-CONHEtPh
- 15           Boc-D-4-GPIA-Gly-Arg-CONHEtPh  
             Boc-D-3-PIG-Gly-Arg-CONHEtPh  
             Boc-D-4-PIG-Gly-Arg-CONHEtPh  
             Boc-D-3-GPIG-Gly-Arg-CONHEtPh  
             Boc-D-4-GPIG-Gly-Arg-CONHEtPh
- 20           BnSO<sub>2</sub>-D-Arg-Gly-Arg-CONHEtPh  
             BnSO<sub>2</sub>-D-3-PIA-Gly-Arg-CONHEtPh  
             BnSO<sub>2</sub>-D-4-PIA-Gly-Arg-CONHEtPh  
             BnSO<sub>2</sub>-D-3-GPIA-Gly-Arg-CONHEtPh  
             BnSO<sub>2</sub>-D-4-GPIA-Gly-Arg-CONHEtPh
- 25           BnSO<sub>2</sub>-D-3-PIG-Gly-Arg-CONHEtPh  
             BnSO<sub>2</sub>-D-4-PIG-Gly-Arg-CONHEtPh  
             BnSO<sub>2</sub>-D-3-GPIG-Gly-Arg-CONHEtPh  
             BnSO<sub>2</sub>-D-4-GPIG-Gly-Arg-CONHEtPh  
             D-Arg-Gly-Arg-CONHEtPh
- 30           D-3-PIA-Gly-Arg-CONHEtPh  
             D-4-PIA-Gly-Arg-CONHEtPh  
             D-3-GPIA-Gly-Arg-CONHEtPh  
             D-4-GPIA-Gly-Arg-CONHEtPh  
             D-3-PIG-Gly-Arg-CONHEtPh
- 35           D-4-PIG-Gly-Arg-CONHEtPh  
             D-3-GPIG-Gly-Arg-CONHEtPh  
             D-4-GPIG-Gly-Arg-CONHEtPh  
             BnSO<sub>2</sub>-D-Arg-Gly-Arg-CON(CH<sub>2</sub>)<sub>4</sub>  
             BnSO<sub>2</sub>-D-3-PIA-Gly-Arg-CON(CH<sub>2</sub>)<sub>4</sub>
- 40

- 5            BnSO<sub>2</sub>-D-4-PIA-Gly-Arg-CON(CH<sub>2</sub>)<sub>4</sub>  
             BnSO<sub>2</sub>-D-3-GPIA-Gly-Arg-CON(CH<sub>2</sub>)<sub>4</sub>  
             BnSO<sub>2</sub>-D-4-GPIA-Gly-Arg-CON(CH<sub>2</sub>)<sub>4</sub>  
             BnSO<sub>2</sub>-D-3-PIG-Gly-Arg-CON(CH<sub>2</sub>)<sub>4</sub>  
             BnSO<sub>2</sub>-D-4-PIG-Gly-Arg-CON(CH<sub>2</sub>)<sub>4</sub>
- 10           BnSO<sub>2</sub>-D-3-GPIG-Gly-Arg-CON(CH<sub>2</sub>)<sub>4</sub>  
             BnSO<sub>2</sub>-D-4-GPIG-Gly-Arg-CON(CH<sub>2</sub>)<sub>4</sub>  
             Boc-D-Arg-Gly-Arg-CONHBn  
             Boc-D-3-PIA-Gly-Arg-CONHBn  
             Boc-D-4-PIA-Gly-Arg-CONHBn
- 15           Boc-D-3-GPIA-Gly-Arg-CONHBn  
             Boc-D-4-GPIA-Gly-Arg-CONHBn  
             Boc-D-3-PIG-Gly-Arg-CONHBn  
             Boc-D-4-PIG-Gly-Arg-CONHBn  
             Boc-D-3-GPIG-Gly-Arg-CONHBn
- 20           Boc-D-4-GPIG-Gly-Arg-CONHBn  
             Boc-D-Arg-Gly-Arg-CONHCH<sub>3</sub>  
             Boc-D-3-PIA-Gly-Arg-CONHCH<sub>3</sub>  
             Boc-D-4-PIA-Gly-Arg-CONHCH<sub>3</sub>  
             Boc-D-3-GPIA-Gly-Arg-CONHCH<sub>3</sub>
- 25           Boc-D-4-GPIA-Gly-Arg-CONHCH<sub>3</sub>  
             Boc-D-3-PIG-Gly-Arg-CONHCH<sub>3</sub>  
             Boc-D-4-PIG-Gly-Arg-CONHCH<sub>3</sub>  
             Boc-D-3-GPIG-Gly-Arg-CONHCH<sub>3</sub>  
             Boc-D-4-GPIG-Gly-Arg-CONHCH<sub>3</sub>
- 30           D-Arg-Gly-Arg-CONHPh  
             Boc-D-3-PIA-Gly-Arg-CONHPh  
             Boc-D-4-PIA-Gly-Arg-CONHPh  
             Boc-D-3-GPIA-Gly-Arg-CONHPh  
             Boc-D-4-GPIA-Gly-Arg-CONHPh
- 35           Boc-D-3-PIG-Gly-Arg-CONHPh  
             Boc-D-4-PIG-Gly-Arg-CONHPh  
             Boc-D-3-GPIG-Gly-Arg-CONHPh  
             Boc-D-4-GPIG-Gly-Arg-CONHPh  
             BnSO<sub>2</sub>-D-Arg-Gly-Arg-CONBn

- 5           BnSO<sub>2</sub>-D-3-PIA-Gly-Arg-CONBn  
            BnSO<sub>2</sub>-D-4-PIA-Gly-Arg-CONBn  
            BnSO<sub>2</sub>-D-3-GPIA-Gly-Arg-CONBn  
            BnSO<sub>2</sub>-D-4-GPIA-Gly-Arg-CONBn  
            BnSO<sub>2</sub>-D-3-PIG-Gly-Arg-CONBn
- 10           BnSO<sub>2</sub>-D-4-PIG-Gly-Arg-CONBn  
            BnSO<sub>2</sub>-D-3-GPIG-Gly-Arg-CONBn  
            BnSO<sub>2</sub>-D-4-GPIG-Gly-Arg-CONBn
7.       A pharmaceutical composition for preventing or treating a condition in a
- 15       mammal characterized by undesired thrombosis comprising a therapeutically acceptable carrier and a therapeutically effective amount of a compound of claim 1, 2, or 6.
8.       A method for preventing or treating a condition in a mammal characterized
- 20       by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of claim 1, 2, or 6.
9.       The method of claim 8, wherein the condition is selected from the group consisting of:
- 25       acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus,
- 30       coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation such as cardiac or other
- 35       intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve, and conditions requiring the fitting of prosthetic devices.
10.      A method for inhibiting the coagulation of biological samples, comprising the administration of a compound of claim 1, 2 or 6.



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 5/06, A61K 38/55</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 96/40743</b>  <b>(43) International Publication Date:</b> 19 December 1996 (19.12.96)
<b>(21) International Application Number:</b> PCT/US96/09285 <b>(22) International Filing Date:</b> 5 June 1996 (05.06.96)  <b>(30) Priority Data:</b> 08/483,470      7 June 1995 (07.06.95)      US  <b>(60) Parent Application or Grant</b> (63) Related by Continuation US      08/483,470 (CIP) Filed on      7 June 1995 (07.06.95)  <b>(71) Applicant (for all designated States except US):</b> COR THER- APEUTICS, INC. [US/US]; 256 East Grand Avenue, South San Francisco, CA 94080 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> MARLOWE, Charles, K. [US/US]; 50 Sequoia Court, San Carlos, CA 94070 (US). SCARBOROUGH, Robert, M. [US/US]; 2544 Belmont Canyon Road, Belmont, CA 94002 (US). LAIBELMAN, Alan, M. [US/US]; 427 Waverley, Menlo Park, CA 94025 (US). SINHA, Uma [IN/US]; 808 Junipero Serra Boulevard, San Francisco, CA 94127 (US). ZHU, Bing-Yan [CA/US]; 3325 Adelaide Way, Belmont, CA 94002-1223 (US).	<b>(74) Agents:</b> MENDLEIN, John, D. et al.; Cooley Godward Castro Huddleson & Tatum, Five Palo Alto Square, 3000 El Camino Real, Palo Alto, CA 94306-2155 (US).  <b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>  <b>(88) Date of publication of the international search report:</b> 23 January 1997 (23.01.97)	
<b>(54) Title:</b> INHIBITORS OF FACTOR Xa  <b>(57) Abstract</b>  Novel compounds, their salts and compositions related thereto having activity against mammalian factor Xa are disclosed. The compounds are useful <i>in vitro</i> or <i>in vivo</i> for preventing or treating coagulation disorders.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam



## INTERNATIONAL SEARCH REPORT

Internat Application No

PCT/US 96/09285

A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 6 C07K5/06 A61K38/55

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO 96 19493 A (CORVAS INTERNATIONAL, INC.) 27 June 1996 see the whole document	1-10
P,X	EP 0 672 659 A (ELI LILLY AND CO.) 20 September 1995 see page 33, line 15 - page 37, line 33	1-10
X	WO 95 09634 A (DU PONT MERCK PHARMACEUTICAL COMPANY ) 13 April 1995 see the whole document	1-10
X	EP 0 417 721 A (MERRELL DOW PHARMACEUTICALS) 20 March 1991 see claims 13-20	1-10
	---	
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*&\* document member of the same patent family

Date of the actual completion of the international search

21 November 1996

Date of mailing of the international search report

0 6. 12. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax (+ 31-70) 340-3016

Authorized officer

Masturzo, P

## INTERNATIONAL SEARCH REPORT

Internati. Application No

PCT/US 96/09285

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 410 411 A (MERRELL DOW PHARMACEUTICALS INC.) 30 January 1991 see claim 18 ---	1-10
X	EP 0 364 344 A (MERRELL DOW PHARMACEUTICAL INC. ) 18 April 1990 see claims 16,17 ---	1-10
X	EP 0 275 101 A (MERRELL DOW PHARMACEUTICALS INC.) 20 July 1988 see the whole document ---	1-10
X	EP 0 195 212 A (MERRELL DOW PHARMACEUTICALS INC.) 24 September 1986 see the whole document ---	1-10
X	US 3 867 364 A (UMEZAWA ET AL.) 18 February 1975 see the whole document ---	1-10
X	CHEMICAL ABSTRACTS, vol. 104, no. 23, 9 June 1986 Columbus, Ohio, US; abstract no. 205517m, "manufacture of a new physiologically active peptide " page 602; XP002019124 see abstract & JP 06 028 990 A (SUNTORY LTD.) 14 February 1985 -----	1-10

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/09285

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Please see Further Information sheet enclosed.
2. ☒ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
Please also see Further Information sheet enclosed.
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 96/ 09285

FURTHER INFORMATION CONTINUED FROM PCT/SA/210

Remark : As the general formula of claim 1 contains no fixed invariable elements, a complete search was considered not to be possible for economical reasons. The searched subject matter includes all the compounds of claim 6 ; this claim includes all the real examples.

Claims completely searched : 6

Claims incompletely searched : 1-5, 7-10

Remark : Although claim 7-10 refer to a method of treatment, the search was carried out and based on the alleged effects of the compounds.

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/US 96/09285

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9619493	27-06-96	AU-A- 4608696	10-07-96
EP-A-672659	20-09-95	CA-A- 2143536 JP-A- 7258287	05-09-95 09-10-95
WO-A-9509634	13-04-95	AU-A- 7966494 EP-A- 0724446 ZA-A- 9407875	01-05-95 07-08-96 09-04-96
EP-A-417721	20-03-91	AT-T- 127474 AU-B- 635961 AU-A- 6223190 CA-A- 2024661 CN-A- 1050198 DE-D- 69022150 JP-A- 3118357	15-09-95 08-04-93 14-03-91 12-03-91 27-03-91 12-10-95 20-05-91
EP-A-410411	30-01-91	AU-B- 632836 AU-A- 5974290 CA-A- 2021660 CN-A- 1049016 FI-B- 94420 HU-B- 209931 IL-A- 95168 JP-A- 3086852	14-01-93 31-01-91 27-01-91 06-02-91 31-05-95 28-12-94 31-10-95 11-04-91
EP-A-364344	18-04-90	AU-B- 617875 AU-A- 4249189 AU-B- 626918 AU-A- 4262589 CA-A- 2000340 CA-A- 2000342 EP-A- 0363284 JP-A- 2256654 JP-A- 2134398 PT-B- 91926 PT-B- 91927	05-12-91 28-06-90 13-08-92 12-04-90 07-04-90 07-04-90 11-04-90 17-10-90 23-05-90 03-07-95 03-07-95
EP-A-275101	20-07-88	AU-B- 614272 AU-A- 1025588	29-08-91 21-07-88

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US 96/09285

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-275101		IL-A- 85105	07-10-94
		JP-A- 63239255	05-10-88
		ZA-A- 8800285	05-07-88
-----			
EP-A-195212	24-09-86	AU-B- 600226	09-08-90
		AU-A- 5288186	07-08-86
		DE-D- 3689314	05-01-94
		DE-T- 3689314	19-05-94
		FI-B- 94254	28-04-95
		FI-C- 94254	10-08-95
		IE-B- 60582	27-07-94
		JP-B- 2529825	04-09-96
		JP-A- 61183253	15-08-86
		US-A- 5496927	05-03-96
-----			
US-A-3867364	18-02-75	BE-A- 742067	04-05-70
		CH-A- 534131	28-02-73
		DE-A- 1958383	04-06-70
		DE-A- 2024330	26-11-70
		FR-A- 2023831	21-08-70
		GB-A- 1292846	11-10-72
		NL-A,B,C 6917470	25-05-70
		SE-B- 369074	05-08-74
-----			